

| Mutual of Omaha Rx Value Plan |

Drugs That Require Prior Authorization (PA) Before Being Approved for Coverage

You will need authorization from **Mutual of Omaha RxSM** (PDP) before filling prescriptions for the drugs shown in the following charts. Mutual of Omaha Rx will only provide coverage after it determines that the drug is being prescribed according to the criteria specified in the chart.

Express Scripts, a pharmacy benefit manager, administers the review process for Mutual of Omaha Rx. To request a prior authorization, please have your physician visit their online portal at **esrx.com/PA**. You, your appointed representative or your prescriber can also request prior authorization by calling Express Scripts toll free at **1.844.374.7377**, 24 hours a day, 7 days a week. Customer Service is available in English and other languages. TTY users should call **1.800.716.3231**.

The formulary may change at any time. You will receive notice when necessary.

Express Scripts is the pharmacy benefit manager for Mutual of Omaha Rx and will be providing some services on behalf of Mutual of Omaha Rx.

Mutual of Omaha Rx (PDP) is a prescription drug plan with a Medicare contract.
Enrollment in the Mutual of Omaha Rx plan depends on contract renewal.

No changes made since 8/25/2020

ACYCLOVIR (TOPICAL)

Products Affected

- acyclovir topical ointment

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Medication history
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ADEMPAS

Products Affected

- Adempas

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	PAH and CTEPH- must be prescribed by or in consultation with a cardiologist or a pulmonologist.
Coverage Duration	1 year
Other Criteria	For PAH - must have PAH (WHO Group 1) and had a right heart catheterization to confirm the diagnosis of PAH (WHO Group 1).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

AFINITOR

Products Affected

- Afinitor
- Afinitor Disperz oral tablet for suspension 2 mg, 3 mg, 5 mg
- everolimus (antineoplastic)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Breast Cancer-HER2 status, hormone receptor (HR) status.
Age Restrictions	Relapsed or refractory classical Hodgkin lymphoma-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.

PA Criteria	Criteria Details
Other Criteria	<p>Breast Cancer-approve if the patient meets ALL the following criteria (A, B, C, D, E, and F):A)patient has recurrent or Stage IV, hormone receptor positive (HR+) [i.e., estrogen receptor positive (ER+) and/or progesterone receptor positive (PR+)] disease AND B)patient has human epidermal growth factor receptor 2 (HER2)-negative breast cancer AND C)patient has tried at least one prior endocrine therapy (e.g., anastrozole, letrozole, or tamoxifen) AND D)patient meets ONE of the following conditions (i or ii): i.patient is a postmenopausal female or a male OR ii.patient is premenopausal or perimenopausal AND is receiving ovarian suppression/ablation with a gonadotropin-releasing hormone (GnRH) agonist (e.g., Lupron [leuprolide], Trelstar [triptorelin], Zoladex (goserelin)), or has had surgical bilateral oophorectomy or ovarian irradiation AND E)patient meets ONE of the following conditions (i or ii):i. If patient is a male AND if everolimus will be used in combination with exemestane, the patient is receiving a gonadotropin-releasing hormone (GnRH) agonist (e.g., Lupron [leuprolide], Trelstar [triptorelin], Zoladex (goserelin)) OR ii. everolimus will be used in combination with exemestane, Faslodex (fulvestrant intramuscular), or tamoxifen AND F) The patient has not had disease progression while on everolimus. Renal Cell Carcinoma (Clear Cell or Non-clear cell histology)-approve if the patient has relapsed or Stage IV disease and if using for clear cell disease, the patient has tried one prior systemic therapy (e.g., Inlyta, Votrient, Sutent, Cabometyx, Nexavar).Tuberous sclerosis complex (TSC) Associated subependymal giant cell astrocytoma (SEGA)-approve if the patient requires therapeutic intervention but cannot be curatively resected. Thymomas and Thymic Carcinomas-approve if the patient has tried one prior chemotherapy (e.g., cisplatin plus doxorubicin, cisplatin plus etoposide, carboplatin plus paclitaxel). TSC associated renal angiomyolipoma -approve. WM/LPL - approve if patient has progressive or relapsed disease or if the patient has not responded to ONE primary therapy (e.g., Velcade with dexamethasone with or without Rituxan, Treanda with Rituxan, Rituxan with cyclophosphamide and dexamethasone, Treanda, Velcade with or</p>

PA Criteria	Criteria Details
	without Rituxan, Velcade with dexamethasone, Kyprolis with Rituxan and dexamethasone, Imbruvica Rituxan). Differentiated (i.e. papillary, follicular, and Hurthle cell) Thyroid Carcinoma-approve if the patient is refractory to radioactive iodine therapy. Endometrial Carcinoma-approve if everolimus will be used in combination with letrozole and the patient has recurrent, metastatic or high-risk disease. GIST-approve if the patient has tried TWO of the following drugs: Sutent, Stivarga, or imatinib AND there is confirmation that everolimus will be used in combination with one of these drugs (Sutent, Stivarga, or imatinib) in the treatment of GIST. Tuberous sclerosis complex (TSC)-associated partial-onset seizures-approve.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Advanced, unresectable or metastatic neuroendocrine tumors of the thymus (Carcinoid tumors). Perivascular Epithelioid Cell Tumors (PEComa), Recurrent Angiomyolipoma, Lymphangioleiomyomatosis, relapsed or refractory classical Hodgkin lymphoma, Waldenstrom's Macroglobulinemia/Lymphoplasmacytic Lymphoma (WM/LPL), Thymomas and Thymic carcinomas, Differentiated (i.e. papillary, follicular, and Hurthle cell) Thyroid Carcinoma, Endometrial Carcinoma, Gastrointestinal Stromal Tumors (GIST) and Recurrent or progressive Meningioma, men with breast cancer

AIMOVIG

Products Affected

- Aimovig Autoinjector

PA Criteria	Criteria Details
Exclusion Criteria	Combination therapy with Ajovy or Emgality
Required Medical Information	Diagnosis, number of migraine headaches per month, prior therapies tried
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Approve if the patient meets the following criteria (A and B): A) Patient has greater than or equal to 4 migraine headache days per month (prior to initiating a migraine-preventative medication), AND B) Patient has tried at least two standard prophylactic pharmacologic therapies, at least one drug each from a different pharmacologic class (e.g., anticonvulsant, beta-blocker), and has had inadequate responses to those therapies or the patient has a contraindication to other prophylactic pharmacologic therapies according to the prescribing physician.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ALECENSA

Products Affected

- Alecensa

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	metastatic NSCLC - is anaplastic lymphoma kinase (ALK)-positive as detected by an approved test.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ALUNBRIG

Products Affected

- Alunbrig oral tablet 180 mg, 30 mg, 90 mg
- Alunbrig oral tablets,dose pack

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	ALK status
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Metastatic NSCLC, must be ALK-positive, as detected by an approved test.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

AMPYRA

Products Affected

- dalfampridine

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	MS. If prescribed by, or in consultation with, a neurologist or MS specialist.
Coverage Duration	Authorization will be for 1 year.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ANABOLIC STEROIDS

Products Affected

- Anadrol-50
- oxandrolone oral tablet 10 mg, 2.5 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	N/A
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients w/Turner's Syndrome or Ullrich-Turner Syndrome (oxandrolone only), management of protein catabolism w/burns or burn injury (oxandrolone only), AIDS wasting and cachexia

ANTIFUNGALS (IV)

Products Affected

- fluconazole in NaCl (iso-osm) intravenous piggyback 200 mg/100 mL, 400 mg/200 mL
- voriconazole intravenous

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

APOKYN

Products Affected

- APOKYN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ARCALYST

Products Affected

- Arcalyst

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent biologic therapy
Required Medical Information	N/A
Age Restrictions	Initial tx CAPS-Greater than or equal to 12 years of age.
Prescriber Restrictions	Initial tx CAPS-prescribed by, or in consultation with, a rheumatologist, geneticist, allergist/immunologist, or dermatologist.
Coverage Duration	3 mos initial, 3 years cont
Other Criteria	CAPS renewal - approve if the patient has had a response as determined by the prescriber.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ARIKAYCE

Products Affected

- Arikayce

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous medication history
Age Restrictions	MAC-18 years and older
Prescriber Restrictions	MAC-Prescribed by a pulmonologist, infectious disease physician or a physician who specializes in the treatment of MAC lung infections. Cystic fibrosis-prescribed by or in consultation with a pulmonologist or physician who specializes in the treatment of cystic fibrosis.
Coverage Duration	1 year
Other Criteria	MAC Lung disease-approve if the patient has NOT achieved negative sputum cultures for Mycobacterium avium complex after a background multidrug regimen AND Arikayce will be used in conjunction to a background multidrug regimen. Note-a multidrug regimen typically includes a macrolide (azithromycin or clarithromycin), ethambutol and a rifamycin (rifampin or rifabutin). Cystic fibrosis-patient has pseudomonas aeruginosa in culture of the airway.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Cystic fibrosis pseudomonas aeruginosa infection

AYVAKIT

Products Affected

- Ayvakit

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	GIST-approve if the patient has unresectable or metastatic disease AND the tumor is positive for platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

BALVERSA

Products Affected

- Balversa

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapies, test results
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Urothelial Carcinoma, locally advanced or metastatic-approve if the patient has susceptible fibroblast growth factor receptor 3 or fibroblast growth factor receptor 2 genetic alterations AND the patient has progressed during or following prior platinum-containing chemotherapy or checkpoint inhibitor therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

BANZEL

Products Affected

- Banzel

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Patients 1 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

BENLYSTA

Products Affected

- Benlysta subcutaneous

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with other biologics or with cyclophosphamide intravenous (IV)
Required Medical Information	Diagnosis, medications that will be used in combination, autoantibody status
Age Restrictions	18 years and older (initial).
Prescriber Restrictions	Prescribed by or in consultation with a rheumatologist, clinical immunologist, nephrologist, neurologist or dermatologist (initial and continuation)
Coverage Duration	Initial-4 months, cont-3 years
Other Criteria	Initial-The patient has autoantibody-positive SLE (i.e., positive for antinuclear antibodies [ANA] and/or anti-double-stranded DNA antibody [anti-dsDNA]) AND Benlysta is being used concurrently with at least one other standard therapy (i.e., antimalarials [e.g., hydroxychloroquine], a systemic corticosteroid [e.g., prednisone], and/or other immunosuppressants [e.g., azathioprine, mycophenolate mofetil, methotrexate]) unless the patient is determined to be intolerant due to a significant toxicity, as determined by the prescribing physician. Continuation-Benlysta is being used concurrently with at least one other standard therapy (i.e., antimalarials [e.g., hydroxychloroquine], a systemic corticosteroid [e.g., prednisone], and/or other immunosuppressants [e.g., azathioprine, mycophenolate mofetil, methotrexate]) unless the patient is determined to be intolerant due to a significant toxicity, as determined by the prescribing physician AND The patient has responded to Benlysta as determined by the prescriber.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

BETASERON

Products Affected

- Betaseron subcutaneous kit

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with other disease-modifying agent used for multiple sclerosis
Required Medical Information	Multiple Sclerosis (MS) diagnosis worded or described as patients with a diagnosis of MS or have experienced an attack and who are at risk of MS.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or after consultation with a neurologist or an MS specialist.
Coverage Duration	Authorization will be for 3 years.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

BOSULIF

Products Affected

- Bosulif oral tablet 100 mg, 400 mg, 500 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Bosulif is being used. For chronic myelogenous leukemia (CML)/ALL, the Philadelphia chromosome (Ph) status of the leukemia must be reported. For ALL, prior therapies tried must be reported to confirm resistance or intolerance.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	For CML, patient must have Ph-positive CML. For ALL, patient must have Ph-positive ALL and has tried ONE other tyrosine kinase inhibitor that is used for Philadelphia chromosome positive ALL (e.g., Gleevec, Sprycel, etc).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients with Philadelphia chromosome positive Acute Lymphoblastic Leukemia

BRAFTOVI

Products Affected

- Braftovi oral capsule 75 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, BRAF V600 status
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Melanoma - approve if the patient has unresectable, advanced or metastatic melanoma AND has a BRAF V600 mutation. Colon or Rectal cancer-approve if the patient meets the following (A, B, and C): A) The patient has BRAF V600E mutation-positive disease AND B) The patient has previously received a chemotherapy regimen for colon or rectal cancer AND C) The agent is prescribed as part of a combination regimen for colon or rectal cancer.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

BRUKINSA

Products Affected

- Brukinsa

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Mantle Cell Lymphoma - approve for 3 years if the patient has tried at least one prior therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

BUPRENORPHINE (SL TABLETS)

Products Affected

- buprenorphine HCl sublingual

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

C1 ESTERASE INHIBITORS

Products Affected

- Cinryze

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	prescribed by or in consultation with an allergist/immunologist or a physician that specializes in the treatment of HAE or related disorders
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

CABLIVI

Products Affected

- Cablivi injection kit

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, concurrent medications
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist
Coverage Duration	3 months
Other Criteria	aTTP-approve if the patient is currently receiving at least one immunosuppressive therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

CABOMETYX

Products Affected

- Cabometyx oral tablet 20 mg, 40 mg, 60 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, histology, RET gene rearrangement status
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Advance Renal Cell Carcinoma (Predominant Clear Cell or Non-Clear Cell Histology)-Approve. Hepatocellular Carcinoma-approve if the patient has been previously treated with at least one other systemic therapy (e.g., Nexavar, Lenvima).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients with Non-Small Cell Lung Cancer with RET Gene Rearrangements

CALQUENCE

Products Affected

- Calquence

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	MCL, CLL and SLL-approve.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

CAPRELSA

Products Affected

- Caprelsa oral tablet 100 mg, 300 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	MTC - approve. DTC - approve if refractory to radioactive iodine therapy.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Differentiated (i.e., papillary, follicular, and Hurthle) Thyroid Carcinoma. Non-Small Cell Lung Cancer with RET Gene Rearrangements

CARBAGLU

Products Affected

- Carbaglu

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a metabolic disease specialist or a specialist who focuses in the treatment of metabolic diseases
Coverage Duration	Pt meets criteria with no genetic test - 3 mo approval. Pt had genetic test - 12 mo approval
Other Criteria	Approve if genetic testing confirmed a mutation leading to N-acetylglutamate synthase deficiency (NAGS) or if the patient has hyperammonemia.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

CAYSTON

Products Affected

- Cayston

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or a physician who specializes in the treatment of cystic fibrosis.
Coverage Duration	1 year
Other Criteria	Approve if the patient has has <i>Pseudomonas aeruginosa</i> in culture of the airway (e.g., sputum culture, oropharyngeal culture, bronchoalveolar lavage culture).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

CHEMET

Products Affected

- Chemet

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Blood lead level
Age Restrictions	Approve in patients between the age of 12 months and 18 years
Prescriber Restrictions	Prescribed by or in consultation with a professional experienced in the use of chelation therapy (eg, a medical toxicologist or a poison control center specialist)
Coverage Duration	Approve for 2 months
Other Criteria	Approve if Chemet is being used to treat acute lead poisoning (not as prophylaxis) and prior to starting Chemet therapy the patient's blood lead level was greater than 45 mcg/dL.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

CHENODAL

Products Affected

- Chenodal

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	For the treatment of gallstones, approve if the patient has tried or is currently using an ursodiol product.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

CHOLBAM

Products Affected

- Cholbam oral capsule 250 mg, 50 mg

PA Criteria	Criteria Details
Exclusion Criteria	Combination Therapy with Chenodal
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with hepatologist, metabolic specialist, or GI
Coverage Duration	3 mos initial, 12 mos cont
Other Criteria	Bile acid synthesis d/o due to SEDs initial - Diagnosis based on an abnormal urinary bile acid as confirmed by Fast Atom Bombardment ionization - Mass Spectrometry (FAB-MS) analysis or molecular genetic testing consistent with the diagnosis. Cont - responded to initial Cholbam tx with an improvement in LFTs AND does not have complete biliary obstruction. Bile-Acid Synthesis Disorders Due to Peroxisomal Disorders (PDs), Including Zellweger Spectrum Disorders initial - PD with an abnormal urinary bile acid analysis by FAB-MS or molecular genetic testing consistent with the diagnosis AND has liver disease, steatorrhea, or complications from decreased fat soluble vitamin absorption (e.g., rickets). Cont - responded to initial Cholbam therapy as per the prescribing physician (e.g., improvements in liver enzymes, improvement in steatorrhea) AND does not have complete biliary obstruction.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

CLOBAZAM

Products Affected

- clobazam oral suspension
- clobazam oral tablet
- Sympazan

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other medications tried
Age Restrictions	2 years and older (initial therapy)
Prescriber Restrictions	Prescribed by or in consultation with a neurologist (initial therapy)
Coverage Duration	1 year
Other Criteria	Lennox-Gastaut Syndrome, initial therapy-patient has tried one of the following: lamotrigine, topiramate, rufinamide, felbamate, or Epidiolex. Treatment refractory seizures/epilepsy, initial therapy-patient has tried and/or is concomitantly receiving at least two other antiepileptic drugs (e.g., valproic acid, lamotrigine, topiramate, clonazepam, levetiracetam, zonisamide, felbamate). Continuation-prescriber confirms patient is responding to therapy.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Dravet Syndrome and treatment-refractory seizures/epilepsy

COMETRIQ

Products Affected

- Cometriq oral capsule 100 mg/day(80 mg x1-20 mg x1), 140 mg/day(80 mg x1-20 mg x3), 60 mg/day (20 mg x 3/day)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	MTC - approve. Non-Small Cell Lung Cancer with RET Gene Rearrangements - approve. Differentiated (i.e., papillary, follicular, and Hurthle) Thyroid Carcinoma-approve if the patient's carcinoma is refractory to radioactive iodine therapy.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Non-Small Cell Lung Cancer with RET Gene Rearrangements, Differentiated (i.e., papillary, follicular, and Hurthle) Thyroid Carcinoma

COPAXONE

Products Affected

- Copaxone subcutaneous syringe 20 mg/mL, 40 mg/mL
- glatiramer subcutaneous syringe 20 mg/mL, 40 mg/mL
- Glatopa subcutaneous syringe 20 mg/mL, 40 mg/mL

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with other disease-modifying agent used for multiple sclerosis
Required Medical Information	Multiple Sclerosis (MS) diagnosis worded or described as patients with a diagnosis of MS or have experienced an attack and who are at risk of MS.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or after consultation with a neurologist or an MS specialist.
Coverage Duration	Authorization will be for 1 year.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

COPIKTRA

Products Affected

- Copiktra

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	CLL/Follicular Lymphoma/SLL/MALT Lymphoma (gastric and non gastric)/marginal zone lymphoma-approve if the patient has tried two prior therapies
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	MALT Lymphoma (gastric and non gastric), Marginal Zone Lymphoma

CORLANOR

Products Affected

- Corlanor oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Previous use of a Beta-blocker, LVEF
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Chronic HF, adults - must have LVEF of less than or equal 35 percent (currently or prior to initiation of Corlanor therapy) AND tried or is currently receiving a Beta-blocker for HF (e.g., metoprolol succinate sustained-release, carvedilol, bisoprolol, carvedilol ER) unless the patient has a contraindication to the use of beta blocker therapy (e.g., bronchospastic disease such as COPD and asthma, severe hypotension or bradycardia). Heart failure due to dilated cardiomyopathy, children-approve.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

COTELLIC

Products Affected

- Cotellic

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Melanoma initial - must have BRAF V600 mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Melanoma (unresectable, advanced or metastatic) - being prescribed in combination with Zelboraf.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

CYSTARAN

Products Affected

- Cystaran

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an ophthalmologist or a metabolic disease specialist or specialist who focuses in the treatment of metabolic diseases
Coverage Duration	1 year
Other Criteria	Approve if the patient has corneal cysteine crystal deposits confirmed by slit-lamp examination
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

DALIRESP

Products Affected

- Daliresp

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Chronic Obstructive Pulmonary Disease (COPD), medications tried.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year.
Other Criteria	COPD, approve in patients who meet all of the following conditions: Patients has severe COPD or very severe COPD, AND Patient has a history of exacerbations, AND Patient has tried a medication from two of the three following drug categories: long-acting beta2-agonist (LABA) [eg, salmeterol, indacaterol], long-acting muscarinic antagonist (LAMA) [eg, tiotropium], inhaled corticosteroid (eg, fluticasone).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

DARAPRIM

Products Affected

- Daraprim
- pyrimethamine

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Patient's immune status
Age Restrictions	N/A
Prescriber Restrictions	Toxoplasma gondii Encephalitis, Chronic Maintenance and Prophylaxis (Primary)-prescribed by or in consultation with an infectious diseases specialist. Toxoplasmosis Treatment-prescribed by or in consultation with an infectious diseases specialist, a maternal-fetal medicine specialist, or an ophthalmologist.
Coverage Duration	12 months
Other Criteria	Toxoplasma gondii Encephalitis, Chronic Maintenance, approve if the patient is immunosuppressed. Toxoplasma gondii Encephalitis Prophylaxis (Primary), approve if the patient is immunosuppressed
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Chronic maintenance and prophylaxis of Toxoplasma Gondii encephalitis

DAURISMO

Products Affected

- Daurismo oral tablet 100 mg, 25 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, medications that will be used in combination, comorbidities
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	AML - approve if Daurismo will be used in combination with cytarabine AND the patient meets i. OR ii: i. patient is using Daurismo for treatment induction and is greater than or equal to 75 years old or the patient has comorbidities that preclude the use of intensive induction chemotherapy according to the prescribing physician, OR ii. patient is continuing Daurismo as post-induction therapy.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients continuing Daurismo as post-induction therapy

DOPTELET

Products Affected

- Doptelet (10 tab pack)
- Doptelet (15 tab pack)
- Doptelet (30 tab pack)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, platelet count, date of procedure
Age Restrictions	18 years and older
Prescriber Restrictions	N/A
Coverage Duration	Thrombo w/chronic liver disease-5 days, chronic ITP-3 years
Other Criteria	Thrombocytopenia with chronic liver disease-Approve if the patient has a current platelet count less than $50 \times 10^9/L$ AND the patient is scheduled to undergo a procedure within 10 to 13 days after starting Doptelet therapy. Chronic ITP-approve if the patient has tried one other therapy or if the patient has undergone splenectomy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

DUPIXENT

Products Affected

- Dupixent Syringe

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with Xolair
Required Medical Information	Diagnosis, prescriber specialty, other medications tried and length of trials
Age Restrictions	AD-6 years and older, asthma-12 years of age and older.
Prescriber Restrictions	Atopic Dermatitis-Prescribed by or in consultation with an allergist, immunologist or dermatologist, asthma-prescribed by or in consultation with an allergist, immunologist or pulmonologist
Coverage Duration	AD-Initial-16 weeks, Cont-1 year, asthma/Rhinosinusitis-initial-6 months, cont 1 year

PA Criteria	Criteria Details
Other Criteria	<p>Atopic Dermatitis-Initial-meets both a and b: a.has used at least one medium, medium-high, high, and/or super-high-potency prescription topical corticosteroid OR has atopic dermatitis affecting ONLY the face, eyes/eyelids, skin folds, and/or genitalia and has tried tacrolimus ointment AND b.Inadequate efficacy was demonstrated with these previously tried topical prescription therapies, according to the prescribing physician.Continuation-Approve if the pt has responded to Dupixent therapy as determined by the prescribing physician. Asthma-Initial-approve if pt meets the following criteria (i, ii, and iii):i.Pt meets ONE of the following criteria (a or b):a)has a blood eosinophil level of greater than or equal to 150 cells per microliter within the previous 6 weeks or within 6 weeks prior to treatment with any anti-interleukin (IL) therapy or Xolair OR b)has oral corticosteroid-dependent asthma, per the prescriber AND ii.has received combination therapy with BOTH of the following (a and b): a)An inhaled corticosteroid (ICS) AND b)At least one additional asthma controller/maintenance medication (NOTE:An exception to the requirement for a trial of one additional asthma controller/maintenance medication can be made if the patient has already received anti-IL-5 therapy or Xolair used concomitantly with an ICS. Use of a combination inhaler containing both an ICS and a LABA would fulfil the requirement for both criteria a and b) AND iii.asthma is uncontrolled or was uncontrolled prior to starting any anti-IL therapy or Xolair as defined by ONE of the following (a, b, c, d or e): a)experienced two or more asthma exacerbations requiring treatment with systemic corticosteroids in the previous year OR b)experienced one or more asthma exacerbation requiring hospitalization or an Emergency Department visit in the previous year OR c)has a forced expiratory volume in 1 second (FEV1) less than 80% predicted OR d)has an FEV1/forced vital capacity (FVC) less than 0.80 OR e)The patient's asthma worsens upon tapering of oral corticosteroid therapy. Continuation-Approve if meets the following criteria (i and ii): i.continues to receive therapy with one inhaled corticosteroid (ICS) or one ICS-containing combination inhaler AND ii.has responded to Dupixent therapy as determined by the prescribing</p>

PA Criteria	Criteria Details
	<p>physician. Chronic rhinosinusitis with Nasal Polyposis-Initial-pt is currently receiving therapy with an intranasal corticosteroid AND is experiencing significant rhinosinusitis symptoms such as nasal obstruction, rhinorrhea, or reduction/loss of smell according to the prescriber AND meets ONE of the following (a or b): a)has received treatment with a systemic corticosteroid within the previous 2 years or has a contraindication to systemic corticosteroid therapy OR b)has had prior surgery for nasal polyps. Continuation-approve if the pt continues to receive therapy with an intranasal corticosteroid AND pt has responded to Dupixent therapy as determined by the prescriber.</p>
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ENBREL

Products Affected

- Enbrel
- Enbrel Mini
- Enbrel SureClick

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with biologic therapy or targeted synthetic DMARD
Required Medical Information	Diagnosis, concurrent medications, previous therapies tried.
Age Restrictions	N/A
Prescriber Restrictions	RA/AS/JIA/JRA,prescribed by or in consult w/ rheumatologist. PsA, prescribed by or in consultation w/ rheumatologist or dermatologist. PP, prescribed by or in consult w/ dermatologist.GVHD,prescribed by or in consult w/ oncologist,hematologist,or physician affiliated w/ transplant center.Behcet's disease,prescribed by or in consult w/ rheumatologist,dermatologist,ophthalmologist,gastroenterologist,or neurologist. Uveitis, prescribed by or in consultation with an ophthalmologist.
Coverage Duration	FDA approved indications - 3 months initial, 3 years cont, others 12 months.

PA Criteria	Criteria Details
Other Criteria	<p>RA initial, patient has tried one conventional synthetic DMARD for at least 3 months (note: patients who have already had a 3-month trial of a biologic for RA are not required to step back and try a conventional synthetic DMARD). JIA/JRA, approve if the pt has aggressive disease, as determined by the prescriber, or the pt has tried one other agent for this condition (eg, MTX, sulfasalazine, leflunomide, NSAID, biologic DMARD or the pt will be started on Enbrel concurrently with MTX, sulfasalazine, or leflunomide or the pt has an absolute contraindication to MTX (eg, pregnancy, breast feeding, alcoholic liver disease, immunodeficiency syndrome, blood dyscrasias), sulfasalazine, or leflunomide. Plaque psoriasis (PP) initial approve if the patient meets one of the following conditions: 1) patient has tried at least one traditional systemic agent for at least 3 months for plaque psoriasis, unless intolerant (eg, MTX, cyclosporine, Soriatane, oral methoxsalen plus PUVA, (note: pts who have already tried a biologic for psoriasis are not required to step back and try a traditional agent first) OR 2) the patient has a contraindication to one oral agent for psoriasis such as MTX. GVHD. Tried or currently is receiving with etanercept 1 conventional GVHD tx (high-dose systemic corticosteroid, CSA, tacrolimus, MM, thalidomide, antithymocyte globulin, etc.). Behcet's. Has tried at least 1 conventional tx (eg, systemic corticosteroid, immunosuppressant, interferon alfa, MM, etc) or adalimumab or infliximab. Uveitis-tried one of the following: periocular, intraocular, or systemic corticosteroid, immunosuppressives, Humira or an infliximab product. RA/AS/JIA/PP/PsA Cont - must have a response to tx according to the prescriber. Behcet's, GVHD, Uveitis Cont-if the patient has had a response to tx according to the prescriber. Clinical criteria incorporated into the Enbrel 25 mg quantity limit edit, approve additional quantity (to allow for 50 mg twice weekly dosing) if one of the following is met: 1) Patient has plaque psoriasis, OR 2) Patient has RA/JIA/PsA/AS and is started and stabilized on 50 mg twice weekly dosing, OR 3) Patient has RA and the dose is being increased to 50 mg twice weekly and patient has taken MTX in combination with Enbrel 50 mg once weekly for at least 2 months, unless MTX is</p>
	<p>contraindicated or intolerant, OR 4) Patient has JIA/PsA/AS and the dose is being increased to 50 mg twice weekly after taking 50 mg once weekly for at least 2 months.</p>
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Graft versus host disease (GVHD), Behcet's disease, Uveitis

EPCLUSA

Products Affected

- Epclusa

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with other direct acting antivirals, excluding ribavirin.
Required Medical Information	Genotype, prescriber specialty, other medications tried or used in combination with requested medication
Age Restrictions	6 years or older
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, infectious diseases physician, or a liver transplant physician
Coverage Duration	Will be c/w AASLD guidance and inclusive of treatment already received for the requested drug
Other Criteria	Criteria will be applied according to AASLD guidelines.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Indications consistent with current AASLD/IDSA guidance

EPIDIOLEX

Products Affected

- Epidiolex

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapies
Age Restrictions	Patients 2 years and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

EPOETIN ALPHA

Products Affected

- Procrit injection solution 10,000 unit/mL, 2,000 unit/mL, 20,000 unit/mL, 3,000 unit/mL, 4,000 unit/mL, 40,000 unit/mL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	CRF anemia in patients not on dialysis. Hemoglobin (Hb) of less than 10.0 g/dL for adults or less than or equal to 11 g/dL for children to start. Hb less than or equal to 11.5 g/dL for adults or 12 g/dL or less for children if previously on epoetin alfa, Mircera or Aranesp. Anemia w/myelosuppressive chemotx.pt must be currently receiving myelosuppressive chemo and Hb 10.0 g/dL or less to start. Hb less than or equal to 12.0 g/dL if previously on epoetin alfa or Aranesp. MDS, approve if Hb is 10 g/dL or less or serum erythropoietin level is 500 mU/mL or less to start. Previously receiving Aranesp or EA, approve if Hb is 12.0 g/dL or less. Anemia in HIV with zidovudine, Hb is 10.0 g/dL or less or endogenous erythropoietin levels are 500 mU/mL or less at tx start. Previously on EA approve if Hb is 12.0 g/dL or less. Surgical pts to reduce RBC transfusions - pt is unwilling or unable to donate autologous blood prior to surgery
Age Restrictions	MDS anemia = 18 years of age and older
Prescriber Restrictions	MDS anemia/myelofibrosis, prescribed by or in consultation with, a hematologist or oncologist.
Coverage Duration	Chemo-6m, Transfus-1m, CKD(dialysis)-3yrs, Myelofibrosis-init-3 mo, cont-1 yr, all others-1 yr
Other Criteria	Myelofibrosis-Initial-patient has a Hb less than 10 or serum erythropoietin less than or equal to 500 Mu/mL. Cont-patient has a Hgb less than or equal to 12 and according to the prescriber the patient has had a response defined as Hb greater than or equal to 10 or an increase of greater than or equal to 2 g/dL.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Anemia due to myelodysplastic syndrome (MDS), Myelofibrosis

ERIVEDGE

Products Affected

- Erivedge

PA Criteria	Criteria Details
Exclusion Criteria	BCC (La or Met) - must not have had disease progression while on Odomzo.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years
Other Criteria	Locally advanced basal cell carcinoma (LABCC), approve if 1. the patient's BCC has recurred following surgery or radiation, OR 2. the patient is not a candidate for surgery and radiation therapy. Central nervous system cancer (this includes brain and spinal cord tumors)- approve if the patient has tried at least one chemotherapy agent and according to the prescriber, the patient has a mutation of the sonic hedgehog pathway.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Central nervous System Cancer

ERLEADA

Products Affected

- Erleada

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ESBRIET

Products Affected

- Esbriet oral capsule
- Esbriet oral tablet 267 mg, 801 mg

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with nintedanib
Required Medical Information	N/A
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	1 year
Other Criteria	IPF - must have FVC greater than or equal to 40 percent of the predicted value AND IPF must be diagnosed with either findings on high-resolution computed tomography (HRCT) indicating usual interstitial pneumonia (UIP) or surgical lung biopsy demonstrating UIP.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

EXJADE

Products Affected

- deferasirox oral tablet, dispersible

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Serum ferritin level
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist
Coverage Duration	1 year
Other Criteria	Transfusion-related chronic iron overload, initial therapy - approve if the patient is receiving blood transfusions at regular intervals for various conditions (eg, thalassemia syndromes, myelodysplastic syndrome, chronic anemia, sickle cell disease) AND prior to starting therapy, the serum ferritin level is greater than 1,000 mcg/L. Non-transfusion-dependent thalassemia syndromes chronic iron overload, initial therapy - approve if prior to starting therapy the serum ferritin level is greater than 300 mcg/L. Continuation therapy - approve if the patient is benefiting from therapy as confirmed by the prescribing physician.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

FARYDAK

Products Affected

- Farydak oral capsule 10 mg, 20 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

FASENRA

Products Affected

- Fasenra

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with Xolair or another Anti-Interleukin (IL) Monoclonal Antibody
Required Medical Information	Diagnosis, severity of disease, peripheral blood eosinophil count, previous therapies tried and current therapies, FEV1/FVC
Age Restrictions	12 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an allergist, immunologist, or pulmonologist
Coverage Duration	Authorization will be for 6 months initial, 12 months continuation.

PA Criteria	Criteria Details
Other Criteria	<p>Initial - must have peripheral blood eosinophil count of greater than or equal to 150 cells per microliter within the previous 6 weeks (prior to treatment with any anti-interleukin (IL)-5 therapy) AND meet both of the following criteria: 1) Patient has received at least 3 consecutive months of combination therapy with an inhaled corticosteroid AND one of the following: inhaled LABA, inhaled long-acting muscarinic antagonist, Leukotriene receptor antagonist, or Theophylline, AND 2) Patient's asthma is uncontrolled or was uncontrolled prior to starting any anti-IL therapy as defined by ONE of the following: a) patient experienced one or more asthma exacerbations requiring treatment with systemic corticosteroids in the previous year, OR b) patient experienced one or more asthma exacerbation requiring hospitalization or an Emergency Department (ED) visit in the previous year, OR c) patient has a FEV1 less than 80 percent predicted, OR d) Patient has an FEV1/FVC less than 0.80, OR e) Patient's asthma worsens upon tapering of oral corticosteroid therapy. NOTE: An exception to the requirement for a trial of one additional asthma controller/maintenance medication can be made if the patient has already received anti-IL-5 therapy (e.g., Cinqair, Fasenra, Nucala) used concomitantly with an ICS for at least 3 consecutive months.</p> <p>Continuation - The patient has responded to Fasenra therapy as determined by the prescribing physician (e.g., decreased asthma exacerbations, decreased asthma symptoms, decreased hospitalizations, emergency department (ED)/urgent care, or physician visits due to asthma, decreased requirement for oral corticosteroid therapy) AND patient continues to receive therapy with an inhaled corticosteroid.</p>
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

FERRIPROX

Products Affected

- Ferriprox oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Serum ferritin level
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist
Coverage Duration	1 year
Other Criteria	Initial therapy - approve if prior to starting therapy the serum ferritin level was greater than 2,500 mcg/L. Continuation therapy - approve if the patient is benefiting from therapy as confirmed by the prescribing physician.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

FIRAZYR

Products Affected

- Firazyr
- icatibant

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by, or in consultation with, an allergist/immunologist or a physician that specializes in the treatment of HAE or related disorders.
Coverage Duration	Authorization will be for 1 year.
Other Criteria	Hereditary Angioedema (HAE) Due to C1 Inhibitor (C1-INH) Deficiency [Type I or Type II] - Treatment of Acute Attacks, Initial Therapy-the patient has HAE type I or type II as confirmed by the following diagnostic criteria (i and ii): i. the patient has low levels of functional C1-INH protein (less than 50% of normal) at baseline, as defined by the laboratory reference values AND ii. the patient has lower than normal serum C4 levels at baseline, as defined by the laboratory reference values. Patients who have treated previous acute HAE attacks with icatibant-the patient has treated previous acute HAE type I or type II attacks with icatibant AND according to the prescribing physician, the patient has had a favorable clinical response (e.g., decrease in the duration of HAE attacks, quick onset of symptom relief, complete resolution of symptoms, decrease in HAE acute attack frequency or severity) with icatibant treatment.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

FIRDAPSE

Products Affected

- Firdapse

PA Criteria	Criteria Details
Exclusion Criteria	History of seizures (initial therapy)
Required Medical Information	Diagnosis, seizure history, lab and test results
Age Restrictions	18 years and older (initial therapy)
Prescriber Restrictions	Prescribed by or in consultation with a neurologist or a neuromuscular specialist (initial therapy)
Coverage Duration	Initial-3 months, Cont-1 year
Other Criteria	Initial therapy-Diagnosis confirmed by at least one electrodiagnostic study (e.g., repetitive nerve stimulation) OR anti-P/Q-type voltage-gated calcium channels (VGCC) antibody testing according to the prescribing physician. Continuation-patient continues to derive benefit (e.g., improved muscle strength, improvements in mobility) from Firdapse, according to the prescribing physician.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

FYCOMPA

Products Affected

- Fycompa oral suspension
- Fycompa oral tablet 10 mg, 12 mg, 2 mg, 4 mg, 6 mg, 8 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Patients 4 years of age or older.
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

GATTEX

Products Affected

- Gattex 30-Vial

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	1 year and older
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist (initial and continuation)
Coverage Duration	1 year
Other Criteria	Initial-approve if the patient is currently receiving parenteral nutrition on 3 or more days per week or according to the prescriber, the patient is unable to receive adequate total parenteral nutrition required for caloric needs. Continuation-approve if the patient has experienced at least a 20 percent decrease from baseline in the weekly volume of parenteral nutrition.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

GILOTRIF

Products Affected

- Gilotrif

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For NSCLC - EGFR exon deletions or mutations or if NSCLC is squamous cell type
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	NSCLC EGFR pos - For the treatment of metastatic non small cell lung cancer (NSCLC) must be used in tumors with non-resistant EGFR mutation positive NSCLC as detected by an approved test. NSCLC metastatic squamous cell must have disease progression with first line treatment with platinum based chemotherapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

GLUCAGON-LIKE PEPTIDE-1 AGONISTS

Products Affected

- Bydureon BCise
- Bydureon subcutaneous pen injector
- Byetta subcutaneous pen injector 10 mcg/dose(250 mcg/mL) 2.4 mL, 5 mcg/dose (250 mcg/mL) 1.2 mL
- Ozempic subcutaneous pen injector 0.25 mg or 0.5 mg(2 mg/1.5 mL), 1 mg/dose (2 mg/1.5 mL)
- Trulicity
- Victoza 3-Pak

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

GONADOTROPIN-RELEASING HORMONE AGONISTS - INJECTABLE LONG ACTING

Products Affected

- Lupron Depot
- Lupron Depot (3 month)
- Lupron Depot (4 month)
- Lupron Depot (6 Month)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	For the treatment of cancer diagnosis must be prescribed by or in consultation with an oncologist.
Coverage Duration	For abnormal uterine bleeding/uterine leiomyomata approve 6 months/all other dx 12 mo
Other Criteria	N/A
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Ovarian cancer, breast cancer, prophylaxis or treatment of uterine bleeding in patients with hematologic malignancy or undergoing cancer treatment or prior to bone marrow/stem cell transplantation, head and neck cancer-salivary gland tumors

GRASTEK

Products Affected

- Grastek

PA Criteria	Criteria Details
Exclusion Criteria	The patient is NOT currently receiving SC or SL allergen immunotherapy
Required Medical Information	Diagnosis
Age Restrictions	5 years through 65 years of age
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	The diagnosis of grass pollen-induced allergic rhinitis must be confirmed by either 1. positive skin test response to a grass pollen from the Pooideae subfamily of grasses (this includes, but is not limited to sweet vernal, Kentucky blue grass, Timothy grass, orchard, or perennial rye grass), or 2. positive in vitro test (i.e., a blood test for allergen-specific IgE antibodies) for a grass in the Pooideae subfamily of grasses. Therapy must be initiated 12 weeks prior to the expected onset of the grass pollen season or therapy is being dosed daily continuously for consecutive grass pollen seasons.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

GROWTH HORMONES

Products Affected

- Norditropin FlexPro subcutaneous pen injector 10 mg/1.5 mL (6.7 mg/mL), 15 mg/1.5 mL (10 mg/mL), 5 mg/1.5 mL (3.3 mg/mL)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	GHD in Children/Adolescents. Pt meets one of the following-1-had 2 GH stim tests with the following-levodopa, insulin-induced hypoglycemia, arginine, clonidine, or glucagon and both are inadequate as defined by a peak GH response which is below the normal reference range of the testing laboratory OR had at least 1 GH test and results show inadequate response and has at least one risk factor for GHD (e.g., ht for age curve deviated down across 2 major height percentiles [e.g., from above the 25 percentile to below the 10 percentile], growth rate is less than the expected normal growth rate based on age and gender, low IGF-1 and/or IGFBP-3 levels). 2.brain radiation or tumor resection and pt has 1 GH stim test and results is inadequate response or has def in at least 1 other pituitary hormone (that is, ACTH, TSH, gonadotropin deficiency [LH and/or FSH] are counted as 1 def], or prolactin).3. congenital hypopituitarism and has one GH stim test with inadequate response OR def in at least one other pituitary hormone and/or the patient has the imaging triad of ectopic posterior pituitary and pituitary hypoplasia with abnormal pituitary stalk 4.pt has panhypopituitarism and has pituitary stalk agenesis, empty sella, sellar or supra-sellar mass lesion, or ectopic posterior pituitary bright spot on MRI or CT or pt has 3 or more pituitary hormone deficiencies or pt has had one GH test and results were inadequate 5.pt had a hypophysectomy. Cont-pt responding to therapy
Age Restrictions	ISS 5 y/o or older, SGA 2 y/o or older, SBS 18 y/o or older
Prescriber Restrictions	GHD (Initial tx children or adolescents w/o hypophysectomy), GHD adults or transitional adolescents, Noonan (initial), Prader Willi (initial for child/adult and cont tx in adults), SHOX (initial), SGA (initial) - prescribed by or in consultation with an endocrinologist. CKD (initial) endocrinologist or nephrologist.
Coverage Duration	ISS - 6 mos intial, 12 months cont tx, SBS-1 month, others 12 mos

PA Criteria	Criteria Details
Other Criteria	<p>GHD initial in adults and adolescents 1. endocrine must certify not being prescribed for anti-aging or to enhance athletic performance, 2. has either childhood onset or adult onset resulting from GHD alone, multiple hormone deficiency from pituitary dx, hypothalamic dz, pituitary surgery, cranial radiation tx, tumor treatment, TBI or subarachnoid hemorrhage, AND 3. meets one of the following - A. childhood onset has known mutations, embryonic lesions, congenital defects or irreversible structural hypothalamic pituitary lesion/damage, B. 3 or more pituitary hormone def (ACTH, TSH, LH/FSH, or prolactin, IGF1 less than 84 mcg/L (Esoterix RIA), AND other causes of low serum IGF-1 have been excluded, C. Neg response to ONE preferred GH stim test (insulin peak response less than or equal to 5 mcg/L, Glucagon peak less than or equal to 3 mcg/L (BMI is less than or equal to 25) or less than or equal to 1 mcg/L (BMI is greater than 25), for transitional adults glucagon peak less than or equal to 3 (BMI is less than 25) or less than or equal to 3 if BMI is greater than or equal to 25 and must also have a second GH stim test with low results, if insulin and glucagon contraindicated then Arginine alone test with peak of less than or equal to 0.4 mcg/L, or Macrilen peak less than 2.8 ng/ml AND BMI is less than or equal to 40 (adults only) AND if a transitional adolescent must be off tx for at least one month before retesting. ISS initial - baseline ht less than the 1.2 percentile or a standard deviation score (SDS) less than -2.25 for age and gender, open epiphyses, does not have CDGP and height velocity is either growth rate (GR) is a. less than 4 cm/yr for pts greater than or equal to 5 or b. growth velocity is less than 10th percentile for age/gender. CKD initial - CKD defined by abnormal CrCl. Noonan initial - baseline height less than 5th percentile. SHOX initial - SHOX def by chromo analysis, open epiphyses, height less than 3rd percentile for age/gender. SGA initial -baseline ht less than 5th percentile for age/gender and born SGA (birth weight/length that is more than 2 SD below mean for gestational age/gender and didn't have sufficient catch up growth by 2-4 y/o). SBS initial pt receiving specialized nutritional support.</p>
	Continuation of tx-approve if the patient has experienced improvement, according to the prescribing physician.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	SHOX, SBS, CKD

HARVONI

Products Affected

- Harvoni oral pellets in packet 33.75-150 mg, 45-200 mg
- Harvoni oral tablet 90-400 mg

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with other direct acting antivirals, excluding ribavirin
Required Medical Information	N/A
Age Restrictions	3 years or older
Prescriber Restrictions	Prescribed by or in consultation w/ GI, hepatologist, ID, or a liver transplant MD
Coverage Duration	Will be c/w AASLD guidance and inclusive of treatment already received for the requested drug
Other Criteria	Criteria will be applied consistent with current AASLD/IDSA guidance.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Indications consistent with current AASLD/IDSA guidance

HETLIOZ

Products Affected

- HetlioZ

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	patient is totally blind with no perception of light
Age Restrictions	18 years or older
Prescriber Restrictions	prescribed by, or in consultation with, a neurologist or a physician who specializes in the treatment of sleep disorders
Coverage Duration	6 mos initial, 12 mos cont
Other Criteria	Initial - dx of Non-24 is confirmed by either assessment of one physiologic circadian phase marker (e.g., measurement of urinary melatonin levels, dim light melatonin onset, assessment of core body temperature), or if assessment of physiologic circadian phase marker cannot be done, the diagnosis must be confirmed by actigraphy plus evaluation of sleep logs. Cont - Approve if pt has achieved adequate results with HetlioZ therapy according to the prescribing physician (e.g., entrainment, clinically meaningful or significant increases in nighttime sleep, clinically meaningful or significant decreases in daytime sleep).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

HIGH RISK MEDICATIONS - BENZODIAZEPINES

Products Affected

- clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg
- diazepam oral concentrate
- diazepam oral solution 5 mg/5 mL (1 mg/mL)
- diazepam oral tablet
- lorazepam oral concentrate
- lorazepam oral tablet 0.5 mg, 1 mg, 2 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
Prescriber Restrictions	N/A
Coverage Duration	Procedure-related sedation = 1mo. All other conditions = 12 months.
Other Criteria	All medically accepted indications other than insomnia, authorize use. Insomnia, may approve lorazepam if the patient has had a trial with two of the following: ramelteon, doxepin 3mg or 6 mg, eszopiclone, zolpidem, or zaleplon. Prior to approval, the physician must have assessed risk versus benefit in prescribing the requested HRM for the patient and must confirm that he/she would still like to initiate/continue therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HIGH RISK MEDICATIONS - BENZTROPINE

Products Affected

- benztropine oral

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For all medically-accepted indications, approve if the prescriber confirms he/she has assessed risk versus benefit in prescribing benztropine for the patient and he/she would still like to initiate/continue therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HIGH RISK MEDICATIONS - CYCLOBENZAPRINE

Products Affected

- cyclobenzaprine oral tablet 10 mg, 5 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Patients aged less than 65 years, approve.
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	The physician has assessed risk versus benefit in using this High Risk Medication (HRM) in this patient and has confirmed that he/she would still like to initiate/continue therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HIGH RISK MEDICATIONS - FIRST GENERATION ANTIHISTAMINES

Products Affected

- hydroxyzine HCl oral tablet
- promethazine oral tablet 25 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months
Other Criteria	For hydroxyzine hydrochloride, authorize use without a previous drug trial for all FDA-approved indications other than anxiety. Approve hydroxyzine hydrochloride if the patient has tried at least two other FDA-approved products for the management of anxiety. Prior to approval, the physician must have assessed risk versus benefit in prescribing the requested HRM for the patient and must confirm that he/she would still like to initiate/continue therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HIGH RISK MEDICATIONS - PHENOBARBITAL

Products Affected

- phenobarbital oral elixir
- phenobarbital oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	Coverage is not provided for use in sedation/insomnia.
Required Medical Information	N/A
Age Restrictions	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For the treatment of seizures, approve only if the patient is currently taking phenobarbital.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HIGH RISK MEDICATIONS - TERTIARY TRICYCLIC ANTIDEPRESSANTS

Products Affected

- amitriptyline
- clomipramine
- doxepin oral capsule
- doxepin oral concentrate
- imipramine HCl
- trimipramine

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	For the treatment of depression, approve if the patient has tried at least two of the following agents: citalopram, escitalopram, fluoxetine, paroxetine, sertraline, venlafaxine, desvenlafaxine, duloxetine, bupropion, mirtazapine or trazodone. For the treatment of pain, may approve amitriptyline (single-entity only, not amitriptyline combination products), or imipramine if the patient has tried at least two of the following agents: duloxetine, pregabalin, gabapentin, venlafaxine or venlafaxine Er. For the treatment of obsessive compulsive disorder (OCD), may approve clomipramine if the patient has tried at least one of the following medications: fluoxetine, fluvoxamine, paroxetine, sertraline, citalopram, escitalopram, or venlafaxine. Prior to approval, the physician must have assessed risk versus benefit in prescribing the requested HRM for the patient and must confirm that he/she would still like to initiate/continue therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HIGH RISK MEDICATIONS- ESTROGENS

Products Affected

- Dotti
- estradiol oral
- estradiol transdermal patch weekly
- norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Previous medication use
Age Restrictions	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months
Other Criteria	For the treatment of Vulvar Vaginal Atrophy, approve if the patient has had a trial of one of the following for vulvar vaginal atrophy (brand or generic): Estradiol Vaginal Cream or estradiol valerate. For prophylaxis of Postmenopausal Osteoporosis, approve if the patient has had a trial of one of the following (brand or generic): alendronate, ibandronate, risidronate or Raloxifene. The physician has assessed risk versus benefit in using this High Risk medication (HRM) in this patient and has confirmed that he/she would still like to initiate/continue therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HUMIRA

Products Affected

- Humira Pen
- Humira Pen Crohns-UC-HS Start
- Humira Pen Psor-Uveits-Adol HS
- Humira subcutaneous syringe kit 10 mg/0.2 mL, 20 mg/0.4 mL, 40 mg/0.8 mL
- Humira(CF) Pedi Crohns Starter subcutaneous syringe kit 80 mg/0.8 mL, 80 mg/0.8 mL-40 mg/0.4 mL
- Humira(CF) Pen Crohns-UC-HS
- Humira(CF) Pen Psor-Uv-Adol HS
- Humira(CF) Pen subcutaneous pen injector kit 40 mg/0.4 mL
- Humira(CF) subcutaneous syringe kit 10 mg/0.1 mL, 20 mg/0.2 mL, 40 mg/0.4 mL

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with another biologic DMARD or targeted synthetic DMARD.
Required Medical Information	Diagnosis, concurrent medications, previous therapies tried
Age Restrictions	Crohn's disease (CD), 6 or older (initial therapy only). Ulcerative colitis (UC), adults.
Prescriber Restrictions	RA/JIA/JRA/Ankylosing spondylitis, prescribed by or in consultation with rheumatologist. Psoriatic arthritis (PsA), prescribed by or in consultation with a rheumatologist or dermatologist. Plaque psoriasis (PP), prescribed by or in consultation with a dermatologist. UC/ CD, prescribed by or in consultation with a gastroenterologist. HS - dermatologist. UV-ophthalmologist
Coverage Duration	initial 3 mo, cont tx 3 years.

PA Criteria	Criteria Details
Other Criteria	<p>RA initial, patient has tried one conventional synthetic DMARD for at least 3 months (note: patients who have already had a 3-month trial of a biologic for RA are not required to step back and try a conventional synthetic DMARD). JIA/JRA initial. Tried another agent (e.g MTX, sulfasalazine, leflunomide, NSAID, or biologic DMARD (eg, etanercept, abatacept, infliximab, anakinra, tocilizumab) or will be starting on adalimumab concurrently with MTX, sulfasalazine, or leflunomide. Approve without trying another agent if pt has absolute contraindication to MTX, sulfasalazine, or leflunomide or if pt has aggressive disease. PP initial, approve if the patient meets one of the following criteria: 1) pt has tried at least one traditional systemic agent (eg, MTX, cyclosporine, acitretin, PUVA) for at least 3 months, unless intolerant (note: pts who have already tried a biologic for psoriasis are not required to step back and try a traditional agent first) OR 2) pt has a contraindication to MTX as determined by the prescribing physician. CD initial. Tried corticosteroids (CSs) or if CSs are contraindicated or if pt currently on CSs or patient has tried one other agent for CD (eg, azathioprine, 6-mercaptopurine, MTX, certolizumab, infliximab, ustekinumab, or vedolizumab) OR pt had ileocolonic resection OR enterocutaneous (perianal or abdominal) or rectovaginal fistulas. UC initial. Patient has tried a systemic therapy (eg, 6-mercaptopurine, azathioprine, CSA, tacrolimus, infliximab, golimumab SC, or a corticosteroid such as prednisone or methylprednisolone) or was intolerant to one of these agents, or the pt has pouchitis and has tried therapy with an antibiotic, probiotic, corticosteroid enema, or mesalamine (Rowasa) enema. FDA approve indications cont tx - must respond to tx as determined by prescriber. HS - tried ONE other therapy (e.g., intralesional or oral corticosteroids, systemic antibiotics, isotretinoin). Clinical criteria incorporated into the Humira 40 mg quantity limit edit allow for approval of additional quantities to accommodate induction dosing. The allowable quantity is dependent upon the induction dosing regimen for the applicable FDA-labeled indications as outlined in product labeling.</p>
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

IBRANCE

Products Affected

- Ibrance

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Breast cancer - approve advanced (metastatic) hormone receptor positive (HR+) [i.e., estrogen receptor positive- (ER+) and/or progesterone receptor positive (PR+)] disease, and HER2-negative breast cancer for patients who have not had disease progression while on Ibrance, Kisqali or Verzenio when the pt meets ONE of the following 1. Pt is postmenopausal and Ibrance will be used in combination with anastrozole, exemestane, or letrozole 2, pt is premenopausal or perimenopausal and is receiving ovarian suppression/ablation with GnRH agonists, or has had surgical bilateral oophorectomy, or ovarian irradiation AND patient meets one of the following critiera: Ibrance will be used in combination with anastrozole, exemestane, or letrozole or Ibrance will be used in combination with Faslodex 3. pt is a man (a man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression) who is receiving GnRH agonist AND Ibrance with be used in combination with anastrozole, exemestane or letrozole or Ibrance will be used in combination with Faslodex 4. Pt is postmenopausal AND Ibrance will be used in combination with Faslodex
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Liposarcoma

ICLUSIG

Products Affected

- Iclusig oral tablet 15 mg, 45 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis the Philadelphia chromosome (Ph) status of the leukemia must be reported. T315I status
Age Restrictions	CML/ALL - Adults
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	CML Ph+, T315I-positive or has tried TWO other TKIs indicated for use in Philadelphia chromosome positive CML (e.g., Gleevec, Sprycel, Tasigna). ALL Ph+, T315I-positive or has tried TWO other TKIs indicated for use in Ph+ ALL (e.g. Gleevec, Sprycel.)
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

IDHIFA

Products Affected

- Idhifa

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	IDH2-mutation status
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	AML - approve if the patient is IDH2-mutation status positive as detected by an approved test
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

IMATINIB

Products Affected

- imatinib oral tablet 100 mg, 400 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis. For indications of CML and ALL, the Philadelphia chromosome (Ph) status of the leukemia must be reported.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	For ALL/CML, new patient must have Ph-positive for approval of imatinib. AIDS related Kaposi's Sarcoma-approve if the patient has tried one prior regimen AND has relapsed or refractory disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Chordoma, advanced or unresectable fibromatosis (desmoid tumors), cKit positive advanced/recurrent or metastatic melanoma, AIDS Related Kaposi's Sarcoma, chronic and pigmented Villonodular Synovitis/Tenosynovial Giant Cell Tumor.

IMBRUVICA

Products Affected

- Imbruvica oral capsule 140 mg, 70 mg
- Imbruvica oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	GVHD-1 year, all others-3 years
Other Criteria	Marginal Zone Lymphoma - Approve. GVHD-Approve if the patient has tried one conventional systemic treatment for graft versus host disease (e.g., corticosteroids [methylprednisolone, prednisone], cyclosporine, tacrolimus, mycophenolate mofetil, imatinib). Diffuse large B-cell lymphoma-approve if the patient is using Imbruvica as second-line or subsequent therapy according to the prescribing physician.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Relapsed or refractory Central Nervous System Lymphoma (Primary). Plus relapsed or refractory Hairy Cell Leukemia. Plus Diffuse Large B-Cell Lymphoma (e.g., follicular lymphoma, gastric MALT lymphoma, nongastric MALT lymphoma, primary DLBCL of the central nervous system).

INCRELEX

Products Affected

- Increlex

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Patients 2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

INJECTABLE TESTOSTERONE PRODUCTS

Products Affected

- testosterone cypionate intramuscular oil 100 mg/mL, 200 mg/mL, 200 mg/mL (1 ML)
- testosterone enanthate

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, lab results
Age Restrictions	Delayed puberty or induction of puberty in males-14 years and older
Prescriber Restrictions	N/A
Coverage Duration	Delayed puberty or induction of puberty in males-6 months, all others-12 months
Other Criteria	Hypogonadism (primary or secondary) in males - initial therapy, approve if all of the following criteria are met: 1) patient has persistent signs and symptoms of androgen deficiency (pre-treatment) [eg, depressed mood, decreased energy, progressive decrease in muscle mass, osteoporosis, loss of libido, AND 2) patient has had two pre-treatment serum testosterone (total or available) measurements, each taken in the morning on two separate days, AND 3) the two serum testosterone levels are both low, as defined by the normal laboratory reference values. Hypogonadism (primary or secondary) in males - continuing therapy, approve if the patient meets all of the following criteria: 1) patient has persistent signs and symptoms of androgen deficiency (pre-treatment) AND 2) patient had at least one pre-treatment serum testosterone level that was low. Delayed puberty or induction of puberty in males - Approve testosterone cypionate or testosterone enanthate. Breast cancer in females. Male is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression. Female is defined as an individual with the biological traits of a woman, regardless of the individual's gender identity or gender expression
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

INLYTA

Products Affected

- Inlyta oral tablet 1 mg, 5 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	Advanced Renal cell carcinoma-approve. Differentiated thyroid cancer, approve if patient is refractory to radioactive iodine therapy.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Differentiated (i.e., papillary, follicular, and Hurthle) Thyroid Carcinoma

INREBIC

Products Affected

- Inrebic

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Myelofibrosis (MF), including Primary MF, Post-Polycythemia Vera MF, and Post-Essential Thrombocythemia MF-approve if the patient has intermediate-2 or high-risk disease.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

IRESSA

Products Affected

- Iressa

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Metastatic NSCLC - The patient has epidermal growth factor receptor (EGFR) exon 19 deletions OR has exon 21 (L858R) substitution mutations as detected by an FDA-approved test.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

IVIG

Products Affected

- Privigen

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	Part B versus D determination per CMS guidance to establish if drug used for PID in pts home.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

JAKAFI

Products Affected

- Jakafi

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	ALL-less than 21 years of age
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	For polycythemia vera patients must have tried hydroxyurea. ALL-approve if the mutation/pathway is Janus associated kinase (JAK)-related. GVHD, chronic-approve if the patient has tried one conventional systemic treatment for graft versus host disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Acute lymphoblastic leukemia, chronic graft versus host disease

KALYDECO

Products Affected

- Kalydeco oral granules in packet
- Kalydeco oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with Orkambi or Symdeko.
Required Medical Information	N/A
Age Restrictions	6 months of age and older
Prescriber Restrictions	prescribed by or in consultation with a pulmonologist or a physician who specializes in CF
Coverage Duration	1 year
Other Criteria	CF - must have one of the following mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene: E56K, P67L, R74W, D110E, D110H, R117C, E193K, L206W, R347H, R352Q, A455E, D579G, S945L, S977F, F1052V, K1060T, A1067T, G1069R, R1070Q, R1070W, F1074L, D1152H, D1270N, G551D, G178R, S549N, S549R, G551S, G1244E, S1251N, S1255P, G1349D, 2789+5G A, 3272-26A G, 3849+10kbC T, 711+3A G, E831X OR R117H AND must NOT be Homozygous for the F508del Mutation in the CFTR Gene or have unknown CFTR gene mutations.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

KISQALI

Products Affected

- Kisqali Femara Co-Pack oral tablet 200 mg/day(200 mg x 1)-2.5 mg, 400 mg/day(200 mg x 2)-2.5 mg, 600 mg/day(200 mg x 3)-2.5 mg
- Kisqali oral tablet 200 mg/day (200 mg x 1), 400 mg/day (200 mg x 2), 600 mg/day (200 mg x 3)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Breast cancer - approve advanced or metastatic hormone receptor positive (HR+) [i.e., estrogen receptor positive (ER+) and/or progesterone receptor positive (PR+)]disease, and HER2-negative breast cancer in patients who have not had disease progression while on Kisqali, Ibrance or Verzenio when the pt meets ONE of the following 1. Pt is postmenopausal and Kisqali will be used in combination with anastrozole, exemestane, or letrozole 2. pt is premenopausal or perimenopausal and is receiving ovarian suppression/ablation with GnRH agonist, or has had surgical bilateral oophorectomy, or ovarian irradiation AND Kisqali will be used in combination with anastrozole, exemestane, or letrozole 3. pt is a man (a man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression) who is receiving GnRH agonist AND Kisqali with be used in combination with anastrozole, exemestane or letrozole. 4. Patient is postmenopausal, pre/perimenopausal or a man and Kisqali (not Co-Pack) will be used in combination with Faslodex 5. Patient is pre/perimenopausal and Kisqali (not Co-Pack) will be used in combination with tamoxifen as first line therapy. If the request is for Kisqali Femara, patients do not need to use in combination with anastrozole, exemestane or letrozole.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Men with breast cancer.

KORLYM

Products Affected

- Korlym

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior surgeries
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist or a physician who specializes in the treatment of Cushing's syndrome.
Coverage Duration	Endogenous Cushing's Synd-1 year. Pt awaiting surgery or response after radiotherapy-4 months
Other Criteria	Endogenous Cushing's Syndrome-Approve if, according to the prescribing physician, the patient is not a candidate for surgery or surgery has not been curative AND if Korlym is being used to control hyperglycemia secondary to hypercortisolism in patients who have type 2 diabetes mellitus or glucose intolerance.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients with Endogenous Cushing's Syndrome, awaiting surgery. Patients with Endogenous Cushing's syndrome, awaiting a response after radiotherapy

KUVAN

Products Affected

- Kuvan

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with Palynziq
Required Medical Information	Diagnosis, Phe concentration
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a specialist who focuses in the treatment of metabolic diseases (initial therapy)
Coverage Duration	Initial-12 weeks, Continuation-1 year
Other Criteria	Initial - approve. Continuation - approve if the patient has had a clinical response (e.g., cognitive and/or behavioral improvements) as determined by the prescribing physician OR patient had a 20% or greater reduction in blood Phe concentration from baseline.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

LENVIMA

Products Affected

- Lenvima oral capsule 10 mg/day (10 mg x 1), 12 mg/day (4 mg x 3), 14 mg/day(10 mg x 1-4 mg x 1), 18 mg/day (10 mg x 1-4 mg x2), 20 mg/day (10 mg x 2), 24 mg/day(10 mg x 2-4 mg x 1), 4 mg, 8 mg/day (4 mg x 2)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	DTC - must be refractory to radioactive iodine treatment for approval. RCC (clear cell or non-clear cell) - approve if the pt meets ALL of the following criteria: 1) pt has relapsed or stage IV disease, 2)if disease is predominant clear-cell histology then the pt has tried one antiangiogenic therapy (eg, Inlyta, Votrient, Sutent, Cabometyx) AND 3)Lenvima will be used in combination with everolimus (Afinitor). MTC-approve if the patient has tried Caprelsa or Cometriq. Anaplastic thyroid cancer-approve if the disease does not have a curative option.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients with Medullary Thyroid Carcinoma (MTC) and anaplastic thyroid carcinoma.

LETAIRIS/TRACLEER

Products Affected

- ambrisentan
- Tracleer

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH) WHO Group 1-results of right heart cath
Age Restrictions	N/A
Prescriber Restrictions	For treatment of pulmonary arterial hypertension, ambrisentan must be prescribed by or in consultation with a cardiologist or a pulmonologist. CTEPH-Tracleer must be prescribed by or in consultation with a cardiologist or pulmonologist
Coverage Duration	Authorization will be for 1 year.
Other Criteria	CTEPH - pt must have tried Adempas, has a contraindication to Adempas, or is currently receiving Tracleer for CTEPH. Pulmonary arterial hypertension (PAH) WHO Group 1, are required to have had a right-heart catheterization to confirm diagnosis of PAH to ensure appropriate medical assessment. Clinical criteria incorporated into the quantity limit edits for sildenafil 20
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Tracleer-Chronic thromboembolic pulmonary hypertension (CTEPH)

LIDOCAINE PATCH

Products Affected

- lidocaine topical adhesive patch,medicated 5 %
- ZTlido

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months
Other Criteria	Chronic back pain-approve if the patient has tried two pharmacologic therapies with each one from a different class of medication used to treat low back pain (e.g., nonsteroidal anti-inflammatory drugs [NSAIDs], muscle relaxants, celecoxib, duloxetine, gabapentin).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Diabetic neuropathic pain, chronic back pain

LONG ACTING OPIOIDS

Products Affected

- buprenorphine transdermal patch weekly 10 mcg/hour, 15 mcg/hour, 20 mcg/hour, 5 mcg/hour
- Butrans transdermal patch weekly 7.5 mcg/hour
- methadone oral solution 10 mg/5 mL, 5 mg/5 mL
- methadone oral tablet 10 mg, 5 mg
- morphine oral tablet extended release
- oxymorphone oral tablet extended release 12 hr

PA Criteria	Criteria Details
Exclusion Criteria	Acute (ie, non-chronic) pain
Required Medical Information	Pain type (chronic vs acute), prior pain medications/therapies tried, concurrent pain medications/therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	For pain severe enough to require daily, around-the-clock, long-term opioid treatment, approve if all of the following criteria are met: 1) patient is not opioid naive, AND 2) non-opioid therapies have been optimized and are being used in conjunction with opioid therapy according to the prescribing physician, AND 3) the prescribing physician has checked the patient's history of controlled substance prescriptions using state prescription drug monitoring program (PDMP), AND 4) the prescribing physician has discussed risks (eg, addiction, overdose) and realistic benefits of opioid therapy with the patient, AND 5) according to the prescriber physician there is a treatment plan (including goals for pain and function) in place and reassessments are scheduled at regular intervals. Patients with cancer, sickle cell disease, in hospice or who reside in a long term care facility are not required to meet above criteria. Clinical criteria incorporated into the quantity limit edits for all oral long-acting opioids require confirmation that the indication is intractable pain (ie, FDA labeled use) prior to reviewing for quantity exception.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off-Label Uses	N/A

LONSURF

Products Affected

- Lonsurf oral tablet 15-6.14 mg, 20-8.19 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	Gastric or Gastroesophageal Junction Adenocarcinoma-approve if the patient has been previously treated with at least two chemotherapy regimens for gastric or gastroesophageal junction adenocarcinoma.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

LORBRENA

Products Affected

- Lorbrena oral tablet 100 mg, 25 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, ALK status, ROS1 status, previous therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	NSCLC - Approve if the patient has ALK-positive metastatic NSCLC and meets one of the following: a) patient has disease progression on Xalkori (crizotinib capsules) and at least one other ALK inhibitor (e.g., Zykadia [ceritinib capsules], Alecensa [alectinib capsules], Alunbrig [brigatinib tablets]), or b) patient has disease progression on Alecensa (alectinib capsules) as the first ALK inhibitor therapy, or c) patient has disease progression on Zykadia (ceritinib capsules) as the first ALK inhibitor therapy. NSCLC-ROS1 Rearrangement-Positive-approve if the patient has disease progression on crizotinib or ceritinib.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Non-small cell lung cancer (NSCLC)-ROS1 Rearrangement-Positive

LYNPARZA

Products Affected

- Lynparza oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years

PA Criteria	Criteria Details
Other Criteria	<p>Ovarian Cancer-Treatment-initial-Approve if the pt meets the following criteria (i and ii): i. The pt has a germline BRCA-mutation as confirmed by an approved test AND per product labeling the pt has progressed on three or more prior lines of chemotherapy.</p> <p>Continuation-approve if the pt has a BRCA mutation (germline) as confirmed by an approved test. Ovarian, Fallopian Tube, or Primary Peritoneal Cancer-Maintenance monotherapy-Approve if the pt meets one of the following criteria (A or B): A) pt meets both of the following criteria for first-line maintenance therapy (i and ii): i. pt has a germline or somatic BRCA mutation-positive disease as confirmed by an approved test AND ii. pt is in complete or partial response to first-line platinum-based chemotherapy regimen (e.g., carboplatin with paclitaxel, carboplatin with doxorubicin, docetaxel with carboplatin) OR B) pt is in complete or partial response after at least two platinum-based chemotherapy regimens (e.g., carboplatin with gemcitabine, carboplatin with paclitaxel, cisplatin with gemcitabine). Ovarian, fallopian tube, or primary peritoneal cancer-maintenance, combination therapy-approve if the medication is used in combination with bevacizumab, the patient has homologous recombination deficiency (HRD)-positive disease, as confirmed by an approved test and the pt is in complete or partial response to first-line platinum-based chemotherapy regimen. Breast Cancer-Approve if the pt meets the following criteria (A, B, C, and D)-A. pt has metastatic, germline BRCA mutation-positive breast cancer AND B. pt has human epidermal growth factor receptor 2 (HER2)-negative breast cancer AND C. The patient meets ONE of the following criteria (i or ii)- i. The patient meets BOTH of the following criteria (a and b)-a) The patient has hormone receptor positive (HR+) [i.e., estrogen receptor positive ER+ and/or progesterone receptor positive PR+] disease AND b) pt meets ONE of the following criteria (1 or 2)-1-pt has been treated with prior endocrine therapy OR-2 pt is considered inappropriate for endocrine therapy OR ii. Pt has triple negative disease (i.e., ER-negative, PR-negative, and HER2-negative) AND D. The patient has been treated with chemotherapy in the neoadjuvant, adjuvant, or</p>

PA Criteria	Criteria Details
	<p>metastatic setting. Pancreatic Cancer-maintenance therapy-approve if the patient has a germline BRCA mutation-positive metastatic disease and the disease has not progressed on at least 16 weeks of treatment with a first-line platinum-based chemotherapy regimen. Prostate cancer-castration resistant-approve if the patient has metastatic disease, the medication is used concurrently with a gonadotropin-releasing hormone (GnRH) analog or the pateint has had a bilateral orchiectomy, the patient has germline or somatic homologous recombination repair (HRR) gene-mutated disease, as confirmed by an approved test, the patient does not have a PPP2R2A mutation and the patient has been previously treated with abiraterone or Xtandi.</p>
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

MEGACE

Products Affected

- megestrol oral suspension 400 mg/10 mL (40 mg/mL), 625 mg/5 mL (125 mg/mL)
- megestrol oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	Coverage is not provided for weight gain for cosmetic reasons.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

MEKINIST

Products Affected

- Mekinist oral tablet 0.5 mg, 2 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Mekinist is being used. For melanoma, thyroid cancer, colon or rectal cancer and NSCLC must have documentation of BRAF V600 mutations
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	Melanoma must be used in patients with BRAF V600 mutation, and patient has unresectable, advanced (including Stage III or Stage IV disease), or metastatic melanoma. Note-This includes adjuvant treatment in patients with Stage III disease with no evidence of disease post-surgery. For NSCLC requires BRAF V600E Mutation and use in combination with Tafenlar. Thyroid cancer, anaplastic-patient has locally advanced or metastatic anaplastic disease AND Mekinist will be taken in combination with Tafenlar, unless intolerant AND the patient has BRAF V600-positive disease. Colon or Rectal cancer-approve if the patient meets the following (A, B, and C): A) The patient has BRAF V600E mutation-positive disease AND B) The patient has previously received a chemotherapy regimen for colon or rectal cancer AND C) The agent is prescribed as part of a combination regimen for colon or rectal cancer.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Colon or rectal cancer

MEKTOVI

Products Affected

- Mektovi

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, BRAF V600 status, concomitant medications
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Melanoma - approve if the patient has unresectable, advanced or metastatic melanoma AND has a BRAF V600 mutation AND Mektovi will be used in combination with Braftovi. Colon or Rectal cancer-approve if the patient meets the following (A, B, and C): A) The patient has BRAF V600E mutation-positive disease AND B) The patient has previously received a chemotherapy regimen for colon or rectal cancer AND C) The agent is prescribed as part of a combination regimen for colon or rectal cancer.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Colon or rectal cancer

MEMANTINE

Products Affected

- memantine oral capsule,sprinkle,ER 24hr
- memantine oral solution
- memantine oral tablet
- memantine oral tablets,dose pack
- Namzaric

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Indication for which memantine is being prescribed.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	N/A
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients with mild to moderate vascular dementia.

MYALEPT

Products Affected

- Myalept

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by, or in consultation with, an endocrinologist or a geneticist physician specialist
Coverage Duration	Authorization will be for 3 years.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

NATPARA

Products Affected

- Natpara

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist.
Coverage Duration	1 year
Other Criteria	Chronic hypoparathyroidism, initial therapy - approve if before starting Natpara, serum calcium concentration is greater than 7.5 mg/dL and 25-hydroxyvitamin D stores are sufficient per the prescribing physician. Chronic hypoparathyroidism, continuing therapy - approve if during Natpara therapy, the patient's 25-hydroxyvitamin D stores are sufficient per the prescribing physician AND the patient is responding to Natapara therapy, as determined by the prescriber.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

NAYZILAM

Products Affected

- Nayzilam

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other medications used at the same time
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	Intermittent Episodes of Frequent Seizure Activity (i.e., seizure clusters, acute repetitive seizures)-approve if the patient is currently receiving maintenance antiepileptic medication(s).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

NERLYNX

Products Affected

- Nerlynx

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Stage of cancer, HER2 status, previous or current medications tried
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Adjuvant tx-12 months, advanced or metastatic dx-3 yrs
Other Criteria	Breast cancer, adjuvant therapy - approve if the patient meets all of the following criteria: Patient has HER2-positive breast cancer AND Patient has completed one year of adjuvant therapy with trastuzumab OR could not tolerate one year of therapy. Breast cancer, advanced or metastatic disease-approve if the patient has HER-2 positive breast cancer, Nerlynx will be used in combination with capecitabine and the patient has tried at least two prior anti-HER2 based regimens in the metastatic setting.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

NEULASTA

Products Affected

- Neulasta subcutaneous syringe

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Cancer patients receiving chemotherapy, if prescribed by or in consultation with an oncologist or hematologist. PBPC - prescribed by or in consultation with an oncologist, hematologist, or physician that specializes in transplantation. Radiation syndrome-prescribed by or in consultation with physician with expertise in treating acute radiation syndrome.
Coverage Duration	Cancer pts receiving chemo-6 mo. PBPC/Radiation Syndrome-1 mo
Other Criteria	Cancer patients receiving chemotherapy, approve if the patient is receiving myelosuppressive anti-cancer medications that are associated with a high risk of febrile neutropenia (the risk is at least 20% based on the chemotherapy regimen), OR the patient is receiving myelosuppressive anti-cancer medications that are associated with a risk of febrile neutropenia but the risk is less than 20% based on the chemotherapy regimen and the patient has one or more risk factors for febrile neutropenia (eg, aged greater than or equal to 65 years), prior chemotherapy or radiation therapy, persistent neutropenia, bone marrow involvement by tumor, recent surgery and/or open wounds, liver and/or renal dysfunction, poor performance status or HIV infection, OR the patient has had a neutropenic complication from prior chemotherapy and did not receive prophylaxis with a colony stimulating factor and a reduced dose or frequency of chemotherapy may compromise treatment.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients undergoing PBPC collection and therapy

NEUPOGEN

Products Affected

- Neupogen

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	AML, HIV/AIDS, MDS - adults
Prescriber Restrictions	Cancer/AML, MDS, ALL, oncologist or a hematologist. Cancer patients receiving BMT and PBPC, prescribed by or in consultation with an oncologist, hematologist, or a physician who specializes in transplantation. SCN, AA - hematologist. HIV/AIDS neutropenia, infectious disease (ID) physician (MD), hematologist, or MD specializing in HIV/AIDS.
Coverage Duration	chemo/SCN/AML-6 mo.HIV/AIDS-4 months.MDS-3 mo.PBPC,Drug induce A/N,AA,ALL,BMT-3 mo.All others=12mo.
Other Criteria	Cancer patients receiving chemotherapy, approve if the patient meets one of the following conditions: patient is receiving myelosuppressive anti-cancer medications that are associated with a high risk of febrile neutropenia (the risk is at least 20% based on the chemotherapy regimen), patient is receiving myelosuppressive anti-cancer medications that are associated with a risk of febrile neutropenia but the risk is less than 20% based on the chemotherapy regimen and the patient has one or more risk factors for febrile neutropenia (eg, aged greater than or equal to 65 years, prior chemotherapy or radiation therapy, persistent neutropenia, bone marrow involvement by tumor, recent surgery and/or open wounds, liver or renal dysfunction, poor performance status, HIV infection), patient has had a neutropenic complication from prior chemotherapy and did not receive prophylaxis with a colony stimulating factor (eg, Leukine, filgrastim products, pegfilgrastim products) and a reduced dose or frequency of chemotherapy may compromise treatment, patient has received chemotherapy has febrile neutropenia and has at least one risk factor (eg, sepsis syndrome, aged greater than 65 years, severe neutropenia [absolute neutrophil count less than 100 cells/mm ³], neutropenia expected to be greater than 10 days in duration, invasive fungal infection).

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Neutropenia associated with human immunodeficiency virus (HIV) or acquired immunodeficiency syndrome (AIDS). Treatment of myelodysplastic syndromes (MDS). Drug induced agranulocytosis or neutropenia. Aplastic anemia (AA). Acute lymphocytic leukemia (ALL).

NEXAVAR

Products Affected

- Nexavar

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	Osteosarcoma, approve if the patient has tried standard chemotherapy and have relapsed/refractory or metastatic disease. GIST, approve if the patient has tried TWO of the following: imatinib mesylate (Gleevec), sunitinib (Sutent), or regorafenib (Stivarga). Differentiated (ie, papillary, follicular, Hurthle) thyroid carcinoma (DTC), approve if the patient is refractory to radioactive iodine treatment. Medullary thyroid carcinoma, approve if the patient has tried vandetanib (Caprelsa) or cabozantinib (Cometriq). AML - Approve if disease is FLT3-ITD mutation positive as detected by an approved test. Renal cell carcinoma (RCC)-approve if the patient has relapsed or Stage IV clear cell histology and the patient has tried at least one prior systemic therapy (e.g., Inlyta, Votrient, Sutent Cabometyx). Ovarian, fallopian tube, primary peritoneal cancer-approve if the patient has platinum resistant disease and Nexavar is used in combination with topotecan.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Osteosarcoma, angiosarcoma, desmoids tumors (aggressive fibromatosis), gastrointestinal stromal tumors (GIST), medullary thyroid carcinoma, Acute Myeloid Leukemia, Chordoma with recurrent disease, solitary fibrous tumor and hemangiopericytoma, ovarian, fallopian tube, primary peritoneal cancer

NINLARO

Products Affected

- Ninlaro

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	MM - be used in combination with Revlimid and dexamethasone OR pt had received at least ONE previous therapy for multiple myeloma (e.g., Thalomid, Revlimid, Pomalyst, Alkeran, dexamethasone, prednisone) OR the agent will be used following autologous stem cell transplantation (ASCT). Systemic light chain amyloidosis-approve if the patient has tried at least one other regimen for this condition. Waldenstrom Macroglobulinemia/lymphoplasmacytic lymphoma-approve if used in combination with a rituximab product and dexamethasone.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients with systemic light chain amyloidosis, Waldenstrom Macroglobulinemia/lymphoplasmacytic lymphoma

NORTHERA

Products Affected

- Northera oral capsule 100 mg, 200 mg, 300 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Medication history
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or a neurologist
Coverage Duration	12 months
Other Criteria	NOH, approve if the patient meets ALL of the following criteria: a) Patient has been diagnosed with symptomatic NOH due to primary autonomic failure (Parkinsons disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, or non-diabetic autonomic neuropathy, AND b) Patient has tried midodrine
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

NUBEQA

Products Affected

- Nubeqa

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

NUEDEXTA

Products Affected

- Nuedexta

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

NUPLAZID

Products Affected

- Nuplazid oral capsule
- Nuplazid oral tablet 10 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

NUVIGIL/PROVIGIL

Products Affected

- modafinil oral tablet 100 mg, 200 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Patients must be greater than or equal to 17 years of age.
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	Excessive sleepiness due to SWSD if the patient is working at least 5 overnight shifts per month. Adjunctive/augmentation treatment for depression in adults if the patient is concurrently receiving other medication therapy for depression.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Excessive daytime sleepiness (EDS) due to myotonic dystrophy - modafinil only. Adjunctive/augmentation for treatment of depression in adults - modafinil only.

OCALIVA

Products Affected

- Ocaliva

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Prescriber specialty, lab values, prior medications used for diagnosis and length of trials
Age Restrictions	18 years and older (initial and continuation therapy)
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant physician (initial and continuation therapy)
Coverage Duration	6 months initial, 3 years cont.
Other Criteria	Initial treatment of PBC-Patient must meet both 1 and 2-1. Patient has a diagnosis of PBC as defined by TWO of the following:a)Alkaline phosphatase (ALP) elevated above the upper limit of normal as defined by normal laboratory reference values b)Positive anti-mitochondrial antibodies (AMAs) c)Histologic evidence of primary biliary cholangitis (PBC) from a liver biopsy 2. Patient meets ONE of the following: a) Patient has been receiving ursodiol therapy for greater than or equal to 1 year and has had an inadequate response. b) Patient is unable to tolerate ursodiol therapy. Cont tx - approve if the patient has responded to Ocaliva therapy as determined by the prescribing physician (e.g., improved biochemical markers of PBC (e.g., alkaline phosphatase [ALP], bilirubin, gamma-glutamyl transpeptidase [GGT], aspartate aminotransferase [AST], alanine aminotransferase [ALT] levels)).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

OCTREOTIDE

Products Affected

- octreotide acetate injection solution

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ODOMZO

Products Affected

- Odomzo

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	BCC - Must not have had disease progression while on Erivedge (vismodegib).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Locally advanced BCC approve if the BCC has recurred following surgery/radiation therapy or if the patient is not a candidate for surgery AND the patient is not a candidate for radiation therapy, according to the prescribing physician. Metastatic BCC - approve.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Metastatic BCC

OFEV

Products Affected

- Ofev

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with pirfenidone
Required Medical Information	N/A
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	1 year
Other Criteria	IPF baseline - must have FVC greater than or equal to 50 percent of the predicted value AND IPF must be diagnosed with either findings on high-resolution computed tomography (HRCT) indicating usual interstitial pneumonia (UIP) or surgical lung biopsy demonstrating UIP. Chronic fibrosing interstitial lung disease-approve if the forced vital capacity is greater than or equal to 45% of the predicted value AND according to the prescriber the patient has fibrosing lung disease impacting more than 10% of lung volume on high-resolution computed tomography AND according to the prescriber the patient has clinical signs of progression.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ORENCIA

Products Affected

- Orenia
- Orenia ClickJect

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with a Biologic DMARD or Targeted Synthetic DMARD.
Required Medical Information	Diagnosis, concurrent medications, previous drugs tried.
Age Restrictions	N/A
Prescriber Restrictions	RA and JIA/JRA prescribed by or in consultation with a rheumatologist. PsA-prescribed by or in consultation with a rheumatologist or dermatologist.
Coverage Duration	3 mos initial, 3 years cont
Other Criteria	RA initial, patient has tried one conventional synthetic DMARD for at least 3 months (note: patients who have already had a 3-month trial of a biologic for RA are not required to step back and try a conventional synthetic DMARD). Juvenile idiopathic arthritis (JIA) [or Juvenile Rheumatoid Arthritis (JRA)], approve if the patient meets one of the following criteria: 1) Patient has tried adalimumab. [Note: the patient does not have to have a trial with Humira if they have had a trial with Enbrel, Actemra IV or infliximab in the past.], OR 2) According to the prescribing physician, the patient has heart failure or a previously treated lymphoproliferative disorder, or a previous serious infection. Cont tx - responded to therapy as per the prescriber.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ORKAMBI

Products Affected

- Orkambi oral granules in packet
- Orkambi oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with Kalydeco or Symdeko
Required Medical Information	N/A
Age Restrictions	2 years of age and older
Prescriber Restrictions	prescribed by or in consultation with a pulmonologist or a physician who specializes in CF
Coverage Duration	3 years
Other Criteria	CF - homozygous for the Phe508del (F508del) mutation in the CFTR gene (meaning the patient has two copies of the Phe508del mutation)
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

OXERVATE

Products Affected

- Oxervate

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by, or in consultation with, an ophthalmologist.
Coverage Duration	2 months
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PALYNZIQ

Products Affected

- Palynziq subcutaneous syringe 10 mg/0.5 mL, 2.5 mg/0.5 mL, 20 mg/mL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, phenylalanine concentrations
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with a specialist who focuses in the treatment of metabolic diseases
Coverage Duration	1 year (initial and continuation)
Other Criteria	Initial therapy - approve if the patient has uncontrolled blood phenylalanine concentrations greater than 600 micromol/L on at least one existing treatment modality (e.g., prior treatment with Kuvan). Maintenance therapy - approve if the patient's blood phenylalanine concentration is less than or equal to 600 micromol/L OR the patient has achieved at least a 20% reduction in blood phenylalanine concentration from pre-treatment baseline.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PEGASYS

Products Affected

- Pegasys ProClick subcutaneous pen injector 180 mcg/0.5 mL
- Pegasys subcutaneous solution
- Pegasys subcutaneous syringe

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous medications tried, liver disease compensation status, concomitant medications for HCV
Age Restrictions	HCV - patients 5 years of age or older, HBV - patients 3 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PEMAZYRE

Products Affected

- Pemazyre

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years
Other Criteria	Cholangiocarcinoma-approve if the patient has unresectable locally advanced or metastatic disease with a fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement, as detected by an approved test AND the patient has been previously treated with at least one systemic therapy regimen
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PHEOCHROMOCYTOMA

Products Affected

- Demser

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior medication trials
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist or a physician who specializes in the management of pheochromocytoma (initial therapy for phenoxybenzamine, initial and continuation therapy for Demser)
Coverage Duration	Authorization will be for 1 year
Other Criteria	If the requested drug is Demser for initial therapy, approve if the patient has tried a selective alpha blocker (e.g., doxazosin, terazosin or prazosin) AND the patient has tried phenoxybenzamine (brand or generic). If the requested drug is Demser for continuation therapy, approve if the patient is currently receiving Demser or has received Demser in the past.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PHOSPHODIESTERASE-5 INHIBITORS FOR PAH

Products Affected

- Alyq
- sildenafil (Pulmonary Arterial Hypertension) oral suspension for reconstitution 10 mg/mL
- sildenafil (Pulmonary Arterial Hypertension) oral tablet 20 mg
- tadalafil (pulm. hypertension)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, right heart cath results
Age Restrictions	N/A
Prescriber Restrictions	For PAH, if prescribed by, or in consultation with, a cardiologist or a pulmonologist.
Coverage Duration	Authorization will be for 1 year.
Other Criteria	Pulmonary arterial hypertension (PAH) WHO Group 1, are required to have had a right-heart catheterization to confirm diagnosis of PAH to ensure appropriate medical assessment. Clinical criteria incorporated into the quantity limit edits for sildenafil 20 mg tablets and suspension require confirmation that the indication is PAH (ie, FDA labeled use) prior to reviewing for quantity exception.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PIQRAY

Products Affected

- Piqray

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Breast Cancer. Approve if the patient meets the following criteria (A, B, C, D, and E): A) The patient is a postmenopausal female, male or premenopausal and is receiving ovarian suppression with a gonadotropin-releasing hormone (GnRH) analog or has had surgical bilateral oophorectomy or ovarian irradiation AND B) The patient has advanced or metastatic hormone receptor (HR)-positive disease AND C) The patient has human epidermal growth factor receptor 2 (HER2)-negative disease AND D) The patient has PIK3CA-mutated breast cancer as detected by an approved test AND E) The patient has progressed on or after at least one prior endocrine-based regimen (e.g., anastrozole, letrozole, exemestane, Faslodex, tamoxifen, toremifene).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

POMALYST

Products Affected

- Pomalyst

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years
Other Criteria	Kaposi Sarcoma-Approve if the patient meets one of the following (i or ii): i. patient is Human Immunodeficiency Virus (HIV)-negative OR ii. patient meets both of the following (a and b): a) The patient is Human Immunodeficiency Virus (HIV)-positive AND b) The patient continues to receive highly active antiretroviral therapy (HAART).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Myelofibrosis, Systemic Light Chain Amyloidosis, Central Nervous System (CNS) Lymphoma, relapsed or refractory disease

PROLASTIN-C

Products Affected

- Prolastin-C intravenous recon soln

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, pretreatment AAT serum concentration
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Alpha1-Antitrypsin (AAT) Deficiency with Emphysema (or COPD) - approve in patients with baseline (pretreatment) AAT serum concentration of less than 80 mg/dL (11 micromol/L).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PROLIA

Products Affected

- Prolia

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other medications for osteoporosis (eg, denosumab [Prolia], bisphosphonates, raloxifene, calcitonin nasal spray [Fortical]), abaloparatide, except calcium and Vitamin D.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 1 year.

PA Criteria	Criteria Details
Other Criteria	<p>Treatment of postmenopausal osteoporosis/Treatment of osteoporosis in men (to increase bone mass) [a man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression], approve if the patient meets one of the following: 1. has had inadequate response after 12 months of therapy with an oral bisphosphonate, had osteoporotic fracture or fragility fracture while receiving an oral bisphosphonate, or intolerability to an oral bisphosphonate, OR 2. the patient cannot take an oral bisphosphonate because they cannot swallow or have difficulty swallowing, they cannot remain in an upright position, or they have a pre-existing GI medical condition, OR 3. pt has tried an IV bisphosphonate (ibandronate or zoledronic acid), OR 4. the patient has severe renal impairment (eg, creatinine clearance less than 35 mL/min) or chronic kidney disease, or if the patient has an osteoporotic fracture or fragility fracture. Treatment of bone loss in patient at high risk for fracture receiving ADT for nonmetastatic prostate cancer, approve if the patient has prostate cancer that is not metastatic to the bone and the patient is receiving ADT (eg, leuprolide, triptorelin, goserelin) or the patient has undergone a bilateral orchiectomy. Treatment of bone loss (to increase bone mass) in patients at high risk for fracture receiving adjuvant AI therapy for breast cancer, approve if the patient has breast cancer that is not metastatic to the bone and in receiving concurrent AI therapy (eg, anastrozole, letrozole, exemestane). Treatment of GIO, approve if pt tried one oral bisphosphonate OR pt cannot take an oral bisphosphonate because the patient cannot swallow or has difficulty swallowing or the patient cannot remain in an upright position post oral bisphosphonate administration or has a pre-existing GI medical condition (eg, patient with esophageal lesions, esophageal ulcers, or abnormalities of the esophagus that delay esophageal emptying [stricture, achalasia]), OR pt has tried zoledronic acid (Reclast), OR pt has severe renal impairment (CrCL less than 35 mL/min) or has CKD or has had an osteoporotic fracture or fragility fracture.</p>
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

PROMACTA

Products Affected

- Promacta oral powder in packet 12.5 mg, 25 mg
- Promacta oral tablet 12.5 mg, 25 mg, 50 mg, 75 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Cause of thrombocytopenia. Thrombocytopenia due to HCV-related cirrhosis, platelet counts. Severe aplastic anemia, platelet counts and prior therapy. MDS-platelet counts.
Age Restrictions	N/A
Prescriber Restrictions	Thrombocytopenia due to chronic ITP or Aplastic Anemia, approve if prescribed by, or after consultation with, a hematologist. Thrombocytopenia due to HCV-related cirrhosis, approve if prescribed by, or after consultation with, a gastroenterologist, hematologist, hepatologist, or a physician who specializes in infectious disease. MDS-presc or after consult with heme/onc.
Coverage Duration	Chronic ITP - 3 years, others 12 months.
Other Criteria	Thrombocytopenia in patients with chronic immune (idiopathic) thrombocytopenia purpura, approve if the patient has tried ONE other therapy or has undergone a splenectomy. Treatment of thrombocytopenia due to HCV-related cirrhosis, approve to allow for initiation of antiviral therapy if the patient has low platelet counts (eg, less than 75,000 mm ³) and the patient has chronic HCV infection and is a candidate for hepatitis C therapy. Aplastic anemia - has low platelet counts at baseline/pretreatment (e.g., less than 30,000 mm ³) AND tried one immunosuppressant therapy (e.g., cyclosporine, mycophenolate mofetil, sirolimus) OR patient will be using Promacta in combination with standard immunosuppressive therapy. MDS-approve if patient has low- to intermediate-risk MDS AND according to the prescriber the patient has clinically-significant thrombocytopenia (e.g., low platelet counts [pretreatment], is platelet transfusion-dependent, active bleeding, and/or a history of bleeding at low platelet counts).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Thrombocytopenia in Myelodysplastic Syndrome (MDS)

QINLOCK

Products Affected

- Qinlock

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years
Other Criteria	Gastrointestinal stromal tumor (GIST), advanced-approve if, according to labeling, the patient has been previously treated with imatinib and at least two other kinase inhibitors, in addition to imatinib.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

QUININE

Products Affected

- quinine sulfate

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

RELISTOR INJECTION

Products Affected

- Relistor subcutaneous solution
- Relistor subcutaneous syringe

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Patients 18 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

REPATHA

Products Affected

- Repatha
- Repatha Pushtronex
- Repatha SureClick

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use of Juxtapid, Kynamro, or Praluent.
Required Medical Information	LDL-C and response to other agents, prior therapies tried, medication adverse event history, medical history
Age Restrictions	ASCVD/HeFH/Primary Hyperlipidemia - 18 yo and older, HoFH 13 yo and older.
Prescriber Restrictions	Prescribed by, or in consultation with, a cardiologist, endocrinologist, or a physician who focuses in the treatment of CV risk management and/or lipid disorders
Coverage Duration	Approve for 3 years for ASCVD/HeFH/HoFH. Approve for 1 year for primary hyperlipidemia.

PA Criteria	Criteria Details
Other Criteria	<p>Hyperlipidemia with HeFH - approve if: 1) diagnosis of HeFH AND 2) tried ONE high intensity statin (i.e. atorvastatin greater than or equal to 40 mg daily or Crestor greater than or equal to 20 mg daily) and LDL remains 70 mg/dL or higher unless pt is statin intolerant defined by experiencing statin related rhabdomyolysis or skeletal-related muscle symptoms while receiving separate trials of atorvastatin and Crestor and during both trials the symptoms resolved upon discontinuation. Hyperlipidemia with ASCVD -approve if: 1) has one of the following conditions: prior MI, h/o ACS, diagnosis of angina, h/o CVA or TIA, PAD, undergone a coronary or other arterial revascularization procedure, AND 2) tried ONE high intensity statin (defined above) and LDL remains 70 mg/dL or higher unless pt is statin intolerant (defined above). HoFH - approve if: 1) has one of the following: a) genetic confirmation of two mutant alleles at the LDLR, APOB, PCSK9, or LDLRAP1 gene locus, OR b) untreated LDL greater than 500 mg/dL (prior to treatment), OR c) treated LDL greater than or equal to 300 mg/dL (after treatment but prior to agents such as Repatha, Kynamro or Juxtapid), OR d) has clinical manifestations of HoFH (e.g., cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas or xanthelasma), AND 2) tried ONE high intensity statin (defined above) for 8 weeks or longer and LDL remains 70 mg/dL or higher unless statin intolerant (defined above). Primary hyperlipidemia (not associated with ASCVD, HeFH, or HoFH)-approve if the patient has tried one high-intensity statin therapy (defined above) and ezetimibe for 8 weeks or longer and LDL remains 100 mg/dL or higher unless statin intolerant (defined above).</p>
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

RETEVMO

Products Affected

- Retevmo

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Medullary Thyroid Cancer/Thyroid Cancer-12 years and older
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years
Other Criteria	Non-Small Cell Lung Cancer (NSCLC)-Approve if the patient has metastatic disease AND the tumor is RET fusion-positive. Medullary Thyroid Cancer-approve if the patient has advanced or metastatic RET-mutant disease and the disease requires treatment with systemic therapy. Thyroid Cancer-approve if the patient has advanced or metastatic RET fusion positive disease, the disease is radioactive iodine-refractory (if radioactive iodine is appropriate) and the disease requires treatment with systemic therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

REVLIMID

Products Affected

- Revlimid

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis and previous therapies or drug regimens tried.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	MCL-approve. MDS-approve if the patient meets one of the following: 1) Pt has symptomatic anemia, OR 2) Pt has transfusion-dependent anemia, OR 3) Pt has anemia that is not controlled with an erythroid stimulating agent (eg, Epogen, Procrit [epoetin alfa injection], Aranesp [darbepoetin alfa injection]). Diffuse, Large B Cell Lymphoma (Non-Hodgkin's Lymphoma)-approve if the pt has tried one other medication treatment regimen. Myelofibrosis-approve if the pt has tried one other therapy. Peripheral T-Cell Lymphoma or T-Cell Leukemia/Lymphoma-approve if the pt has tried one other chemotherapy regimen.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Systemic Amyloidosis Light Chain, Diffuse Large B Cell Lymphoma (Non-Hodgkin's Lymphoma), Myelofibrosis. Castleman's Disease, Hodgkin lymphoma (Classical), Peripheral T-Cell Lymphoma, T-Cell Leukemia/Lymphoma.

RINVOQ

Products Affected

- Rinvoq

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with a biologic or with a targeted synthetic DMARD. Concurrent use with other potent immunosuppressants.
Required Medical Information	Diagnosis, concurrent medications, previous drugs tried.
Age Restrictions	18 years and older
Prescriber Restrictions	RA, prescribed by or in consultation with a rheumatologist.
Coverage Duration	Authorization will be for 3 months initial, 3 years cont.
Other Criteria	RA initial-approve if the patient has tried one conventional synthetic DMARD for at least 3 months (note: patients who have already had a 3-month trial of a biologic for RA are not required to step back and try a conventional synthetic DMARD). Continuation Therapy - Patient must have responded, as determined by the prescriber
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ROZLYTREK

Products Affected

- Rozlytrek oral capsule 100 mg, 200 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Solid Tumors-12 years and older
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Solid Tumors-Approve if the patient meets the following criteria (A, B, and C): A) The patient has locally advanced or metastatic solid tumor AND B) The patient's tumor has neurotrophic receptor tyrosine kinase (NTRK) gene fusion AND C) The patient meets one of the following criteria (i or ii): i. The patient has progressed on prior therapies OR ii. There are no acceptable standard therapies and the medication is used as initial therapy. Non-Small Cell Lung Cancer-Approve if the patient has ROS1-positive metastatic disease.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

RUBRACA

Products Affected

- Rubraca

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Rubraca is being used. BRCA-mutation (germline or somatic) status. Other medications tried for the diagnosis provided
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3years
Other Criteria	Ovarian, Fallopian Tube or Primary Peritoneal Cancer-treatment - Approve if the patient meets the following criteria (i and ii): i.The patient has a BRCA-mutation (germline or somatic) as confirmed by an approved test, AND ii.The patient has progressed on two or more prior lines of chemotherapy. Maintenance Therapy of Ovarian, Fallopian tube or Primary peritoneal cancer-Approve if the patient has recurrent disease and patient is in complete or partial response after at least two lines of platinum-based chemotherapy. Castration-Resistant Prostate Cancer - Approve if the patient meets the following criteria (A, B, C, and D): A) The patient has metastatic disease that is BRCA-mutation positive (germline and/or somatic) AND B) The patient meets one of the following criteria (i or ii): i. The medication is used concurrently with a gonadotropin-releasing hormone (GnRH) analog OR ii. The patient has had a bilateral orchiectomy AND C) The patient has been previously treated with at least one androgen receptor-directed therapy AND D) The patient meets one of the following criteria (i or ii): i. The patient has been previously treated with at least one taxane-based chemotherapy OR ii. The patient is not a candidate or is intolerant to taxane-based chemotherapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

RYDAPT

Products Affected

- Rydapt

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For AML, FLT3 status
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	AML-approve if the patient is FLT3-mutation positive as detected by an approved test AND the patient is receiving Rydapt in one of the following settings (i, ii, iii, or iv)-i. Induction therapy in combination with cytarabine and daunorubicin OR ii. After standard-dose cytarabine induction/reinduction, along with cytarabine and daunorubicin OR iii. Post remission or consolidation therapy in combination with cytarabine OR iv. Maintenance therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

SABRIL

Products Affected

- vigabatrin
- Vigadrone

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, medication history (complex partial seizures)
Age Restrictions	Refractory complex partial seizures - patients 10 years of age or older. Infantile spasms/West Syndrome - patients 1 month to 2 years of age
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

SAMSCA

Products Affected

- Samsca oral tablet 15 mg, 30 mg

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with Jynarque.
Required Medical Information	Serum sodium less than 125 mEq/L at baseline or less marked hyponatremia, defined as serum sodium less than 135 mEq/L at baseline, that is symptomatic (eg, nausea, vomiting, headache, lethargy, confusion).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 30 days
Other Criteria	Hyponatremia - Pt must meet ONE of the following: 1. serum sodium less than 125 mEq/L at baseline, OR 2. marked hyponatremia, defined as less than 135 mEq/L at baseline, that is symptomatic (eg, nausea, vomiting, headache, lethargy, confusion), OR 3. patient has already been started on Samsca and has received less than 30 days of therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

SIGNIFOR

Products Affected

- Signifor

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist or a physician or specializes in the treatment of Cushing's syndrome.
Coverage Duration	Cushing's dis-Initial-4 mo, Cont-1 year. Pt awaiting surgery or response after radiotherapy-4 mo
Other Criteria	Cushing's disease, initial therapy - approve if, according to the prescribing physician, the patient is not a candidate for surgery, or surgery has not been curative. Cushing's disease, continuation therapy - approve if the patient has already been started on Signifor and, according to the prescribing physician, the patient has had a response and continuation of therapy is needed to maintain response.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

SIRTURO

Products Affected

- Sirturo oral tablet 100 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, concomitant therapy
Age Restrictions	Patients 12 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

SKYRIZI

Products Affected

- Skyrizi subcutaneous syringe kit

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent Use with other Biologics or Targeted Synthetic Disease-Modifying Antirheumatic Drugs (DMARDs)
Required Medical Information	Diagnosis, Previous medication use
Age Restrictions	18 years of age and older (initial therapy)
Prescriber Restrictions	PP-Prescribed by or in consultation with a dermatologist (initial therapy)
Coverage Duration	3 mos initial, 3 years cont
Other Criteria	Initial Therapy-The patient meets ONE of the following conditions (a or b): a) The patient has tried at least one traditional systemic agent for psoriasis (e.g., methotrexate [MTX], cyclosporine, acitretin tablets, or psoralen plus ultraviolet A light [PUVA]) for at least 3 months, unless intolerant. NOTE: An exception to the requirement for a trial of one traditional systemic agent for psoriasis can be made if the patient has already had a 3-month trial or previous intolerance to at least one biologic (e.g., an adalimumab product [Humira], a certolizumab pegol product [Cimzia], an etanercept product [Enbrel, Erelzi], an infliximab product [e.g., Remicade, Inflectra, Renflexis], Cosentyx [secukinumab SC injection], Ilumya [tildrakizumab SC injection], Siliq [brodalumab SC injection], Stelara [ustekinumab SC injection], Taltz [ixekizumab SC injection], or Tremfya [guselkumab SC injection]). These patients who have already tried a biologic for psoriasis are not required to 'step back' and try a traditional systemic agent for psoriasis)b) The patient has a contraindication to methotrexate (MTX), as determined by the prescribing physician.Continuation Therapy - Patient must have responded, as determined by the prescriber.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

SOMATULINE DEPOT (INJECTION AND SC INJECTION)

Products Affected

- Somatuline Depot

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous treatments/therapies
Age Restrictions	N/A
Prescriber Restrictions	Acromegaly-prescribed by or in consultation with an endocrinologist. Carcinoid syndrome-prescribed by or in consultation with an oncologist, endocrinologist or gastroenterologist. All neuroendocrine tumors-prescribed by or in consultation with an oncologist, endocrinologist, or gastroenterologist. Pheochromocytoma/paraganglioma-prescribed by or in consultation with an endo/onc/neuro.
Coverage Duration	1 year
Other Criteria	Acromegaly-approve if the patient has a pre-treatment (baseline) insulin-like growth factor-1 (IGF-1) level above the upper limit of normal based on age and gender for the reporting laboratory AND the patient meets i., ii., or iii: i. has had an inadequate response to surgery and/or radiotherapy or ii. is not an appropriate candidate for surgery and/or radiotherapy or iii. the patient is experiencing negative effects due to tumor size (e.g., optic nerve compression).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Pheochromocytoma/paraganglioma

SOMAVERT

Products Affected

- Somavert

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

SPRYCEL

Products Affected

- Sprycel oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Sprycel is being used. For indications of CML and ALL, the Philadelphia chromosome (Ph) status of the leukemia must be reported.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	For CML, new patient must have Ph-positive CML for approval of Sprycel. For ALL, new patient must have Ph-positive ALL for approval of Sprycel. GIST - has D842V mutation AND previously tried Sutent and Gleevec.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	GIST, chondrosarcoma or chordoma

STELARA

Products Affected

- Stelara subcutaneous

PA Criteria	Criteria Details
Exclusion Criteria	Ustekinumab should not be given in combination with a Biologic DMARD or Targeted Synthetic DMARD
Required Medical Information	Diagnosis, concurrent medications, previous drugs tried.
Age Restrictions	Adults-PsA and CD. PP-12 years and older.
Prescriber Restrictions	PP-Prescr/consult w/derm.PsA-prescr/consult w/rheum or derm.CD/UC-prescr/consult w/gastro.
Coverage Duration	PP/PsA Init-3mo,CD/UC load-approve 1 dose IV,CD/UC post IV load-SC 3 mo,cont tx-SC 3 yr
Other Criteria	PP initial - approve Stelara SC.CD, initial therapy - approve 3 months of the SC formulation if the patient meets ONE of the following criteria: 1) patient has tried or is currently taking corticosteroids, or corticosteroids are contraindicated, OR 2) patient has tried one other agent for CD. UC, initial therapy - approve 3 months of the SC formulation if the patient has had a trial of one systemic agent for UC. PP/PsA/CD/UC cont - approve Stelara SC if according to the prescribing physician, the patient has responded to therapy.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

STIVARGA

Products Affected

- Stivarga

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Stivarga is being used. Prior therapies tried. For metastatic CRC, KRAS/NRAS mutation status.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	For GIST, patient must have previously been treated with imatinib (Gleevec) and sunitinib (Sutent). For HCC, patient must have previously been treated with at least one tyrosine kinase inhibitor (e.g., Nexavar, Lenvima). Soft tissue sarcoma-approve if the patient has non-adipocytic extremity/superficial trunk, head/neck or retroperitoneal/intra-abdominal sarcoma OR pleomorphic rhabdomyosarcoma. Osteosarcoma-approve if the patient has relapsed/refractory or metastatic disease and the requested medication is being used as subsequent therapy.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Soft tissue Sarcoma, Osteosarcoma

SUTENT

Products Affected

- Sutent

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	Gastrointestinal stromal tumors (GIST), approve if the patient has tried imatinib (Gleevec). Chordoma, approve if the patient has recurrent disease. Differentiated thyroid carcinoma, approve if the patient is refractory to radioactive iodine therapy. Medullary thyroid carcinoma, approve if the patient has tried vandetanib (Caprelsa) or cabozantinib (Cometriq). Meningioma, approve if the patient has recurrent or progressive disease. Thymic carcinoma - has tried chemotherapy or radiation therapy. Renal Cell Carcinoma (RCC), clear cell or non-clear cell histology-approve if the patient is at high risk of recurrent clear cell RCC following nephrectomy and Sutent is used for adjuvant therapy or if the patient has relapsed or Stage IV disease. Neuroendocrine tumors of the pancreas-approve for advanced or metastatic disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Chordoma, angiosarcoma, solitary fibrous tumor/hemangiopericytoma, alveolar soft part sarcoma (ASPS), differentiated (ie, papillary, follicular, and Hurthle) thyroid carcinoma, medullary thyroid carcinoma, meningioma, thymic carcinoma.

SYLATRON

Products Affected

- Sylatron subcutaneous kit 200 mcg, 300 mcg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, time elapsed since surgical resection including complete lymphadenectomy
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

SYMDEKO

Products Affected

- Symdeko

PA Criteria	Criteria Details
Exclusion Criteria	Patients with unknown CFTR gene mutations
Required Medical Information	Diagnosis, specific CFTR gene mutations
Age Restrictions	Six years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or a physician who specializes in CF
Coverage Duration	3 years
Other Criteria	CF - must have at least one of the following mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene: E56K, P67L, R74W, D110E, D110H, R117C, E193K, L206W, R347H, R352Q, A455E, D579G, 711+3A G, S945L, S977F, F1052V, E831X, K1060T, A1067T, R1070W, F1074L, D1152H, D1270N, 2789+5G A, 3272-26A G, or 3849 + 10kbC T OR the patient has two copies of the F508del mutation
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

SYMLIN

Products Affected

- SymlinPen 120
- SymlinPen 60

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

SYPRINE

Products Affected

- trientine

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, medication history, pregnancy status, disease manifestations
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant physician.
Coverage Duration	Authorization will be for 3 years.
Other Criteria	For Wilson's Disease, approve if the patient meets ONE of the following: 1) Patient has tried a penicillamine product and per the prescribing physician the patient is intolerant to penicillamine therapy, OR 2) Per the prescribing physician, the patient has clinical features indicating the potential for intolerance to penicillamine therapy (ie, history of any renal disease, congestive splenomegaly causing severe thrombocytopenia, autoimmune tendency), OR 3) Per the prescribing physician, the patient has a contraindication to penicillamine therapy, OR 4) The patient has neurologic manifestations of Wilson's disease, OR 5) The patient is pregnant.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

TABRECTA

Products Affected

- Tabrecta

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years
Other Criteria	Non-Small Cell Lung Cancer (NSCLC)-Approve if the patient has metastatic disease AND the tumor is positive for a mutation that leads to mesenchymal-epithelial transition (MET) exon 14 skipping, as detected by an approved test.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

TAFAMIDIS

Products Affected

- Vyndamax

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with Onpattro or Tegsedi. Concurrent use of Vyndaqel and Vyndamax.
Required Medical Information	Diagnosis, genetic tests and lab results
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consult with a cardiologist/physician who specializes in the tx of amyloidosis
Coverage Duration	1 year
Other Criteria	Cardiomyopathy of Wild-Type or Hereditary Transthyretin Amyloidosis-approve if the diagnosis was confirmed by one of the following (i, ii or iii): i. A technetium pyrophosphate scan (i.e., nuclear scintigraphy), ii. Amyloid deposits are identified on cardiac biopsy OR iii. patient had genetic testing which, according to the prescriber, identified a TTR mutation AND Diagnostic cardiac imaging (e.g., echocardiogram, cardiac magnetic imaging) has demonstrated cardiac involvement (e.g., increased thickness of the ventricular wall or interventricular septum).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

TAFINLAR

Products Affected

- Tafinlar

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Tafinlar is being used. BRAF V600 mutations
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	Melanoma with BRAF V600 mutation AND patient has unresectable, advanced (including Stage III or Stage IV disease) or metastatic melanoma. Note-This includes adjuvant treatment in patients with Stage III disease with no evidence of disease post-surgery. For NSCLC, must have BRAF V600E mutation. Thyroid Cancer, anaplastic-must have BRAF V600-positive disease AND Tafinlar will be taken in combination with Mekinist, unless intolerant AND the patient has locally advanced or metastatic anaplastic disease. Thyroid Cancer, differentiated (i.e., papillary, follicular, or Hurthle cell) AND the patient has disease that is refractory to radioactive iodine therapy AND the patient has BRAF-positive disease. Colon or Rectal cancer-approve if the patient meets the following (A, B, and C): A) The patient has BRAF V600E mutation-positive disease AND B) The patient has previously received a chemotherapy regimen for colon or rectal cancer AND C) The agent is prescribed as part of a combination regimen for colon or rectal cancer.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients with Differentiated Thyroid Cancer, Colon or rectal cancer

TAGRISSO

Products Affected

- Tagrisso

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	NSCLC - prior therapies and EGFR T790M mutation or EGFR exon 19 deletion or exon 21 (L858R) substitution
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	NSCLC - Must have metastatic epidermal growth factor receptor (EGFR) T790M mutation-positive NSCLC as detected by an approved test AND has progressed on one of the EGFR tyrosine kinase inhibitors (e.g., Tarceva, Iressa, Vizimpro or Gilotrif) therapy OR Advanced or metastatic Non-Small Cell Lung Cancer (NSCLC) who have EGFR exon 19 deletion or exon 21 (L858R) substitution as detected by an approved test.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

TALZENNA

Products Affected

- Talzenna oral capsule 0.25 mg, 1 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, BRCA mutation status, HER2 status
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Locally-advanced or metastatic breast cancer-approve if the patient has germline BRCA mutation-positive AND human epidermal growth factor receptor 2 (HER2) negative disease
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

TARCEVA

Products Affected

- erlotinib oral tablet 100 mg, 150 mg, 25 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Advanced, recurrent, or metastatic non small cell lung cancer (NSCLC), EGFR mutation or gene amplification status.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	Metastatic NSCLC, approve if the patient meets both of the following: 1. patient is EGFR mutation positive, AND 2. patient has EGFR exon 19 deletions OR exon 21 (L858R) substitution mutations as detected by an FDA-approved test. Advanced RCC, approve if the patient has non-clear cell histology.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Renal Cell Carcinoma and Bone Cancer-Chordoma.

TARGRETIN ORAL

Products Affected

- bexarotene

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapies tried
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist (initial and continuation)
Coverage Duration	3 years
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

TARGRETIN TOPICAL

Products Affected

- Targretin topical

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapies tried
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist (initial and continuation)
Coverage Duration	3 years
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

TASIGNA

Products Affected

- Tasigna oral capsule 150 mg, 200 mg, 50 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Tasigna is being used. For indication of CML and ALL, the Philadelphia chromosome (Ph) status of the leukemia must be reported. For indication of gastrointestinal stromal tumor (GIST) and ALL, prior therapies tried.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	For CML, new patient must have Ph-positive CML for approval of Tasigna. For GIST, patient must have tried TWO of the following - sunitinib (Sutent), imatinib (Gleevec), or regorafenib (Stivarga). For ALL, Approve if the patient has tried one other tyrosine kinase inhibitor that is used for Philadelphia chromosome positive ALL (e.g., Gleevec, Sprycel, etc).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Philadelphia positive Acute Lymphoblastic Leukemia (ALL) and Gastrointestinal Stromal Tumor (GIST)

TAZORAC

Products Affected

- tazarotene
- Tazorac topical cream 0.05 %

PA Criteria	Criteria Details
Exclusion Criteria	Cosmetic uses
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Acne vulgaris after a trial with at least 1 other topical retinoid product (eg, tretinoin cream/gel/solution/microgel, adapalene).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TAZVERIK

Products Affected

- Tazverik

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Epithelioid Sarcoma-16 years and older, Follicular Lymphoma-18 years and older
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Epithelioid Sarcoma-approve if the patient has metastatic or locally advanced disease and the patient is not eligible for complete resection. Follicular Lymphoma-approve if the patient has relapsed or refractory disease and according to the prescriber, there are no appropriate alternative therapies or the patient's tumor is positive for an EZH2 mutation and the patient has tried at least two prior systemic therapies
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

TECFIDERA

Products Affected

- Tecfidera

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with other disease-modifying agents used for multiple sclerosis (MS)
Required Medical Information	MS, patient must have a relapsing form of MS (RRMS, SPMS with relapses, or PRMS).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist or MS specialist.
Coverage Duration	Authorization will be for 1 year.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

TERIPARATIDE

Products Affected

- teriparatide

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other medications for osteoporosis (eg, denosumab [Prolia], bisphosphonates, raloxifene, calcitonin nasal spray [Fortical], abaloparatide), except calcium and Vitamin D.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 2 years of therapy over a patient's lifetime
Other Criteria	Treatment of PMO, approve if pt has tried one oral bisphosphonate OR pt cannot take an oral bisphosphonate because the pt cannot swallow or has difficulty swallowing or the pt cannot remain in an upright position post oral bisphosphonate administration or pt has a pre-existing GI medical condition (eg, patient with esophageal lesions, esophageal ulcers, or abnormalities of the esophagus that delay esophageal emptying [stricture, achalasia]), OR pt has tried an IV bisphosphonate (ibandronate or zoledronic acid), OR pt has severe renal impairment (creatinine clearance less than 35 mL/min) or CKD or pt has had an osteoporotic fracture or fragility fracture. Increase bone mass in men (a man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression) with primary or hypogondal osteoporosis/Treatment of GIO, approve if pt tried one oral bisphosphonate OR pt cannot take an oral bisphosphonate because the patient cannot swallow or has difficulty swallowing or the patient cannot remain in an upright position post oral bisphosphonate administration or has a pre-existing GI medical condition (eg, patient with esophageal lesions, esophageal ulcers, or abnormalities of the esophagus that delay esophageal emptying [stricture, achalasia]), OR pt has tried zoledronic acid (Reclast), OR pt has severe renal impairment (CrCL less than 35 mL/min) or has CKD or has had an osteoporotic fracture or fragility fracture.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

THALOMID

Products Affected

- Thalomid oral capsule 100 mg, 150 mg, 200 mg, 50 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	<p>Discoid lupus erythematosus or cutaneous lupus erythematosus, approve if the patient has tried two other therapies (eg, corticosteroids [oral, topical, intralesional], hydroxychloroquine, tacrolimus [Protopic], methotrexate, dapsone, acitretin [Soriatane]).</p> <p>Myelofibrosis, approve if the patient has tried one other therapy (eg, ruxolitinib [Jakafi], danazol, epoetin alfa [Epogen/Procrit], prednisone, lenalidomide [Revlimid], hydroxyurea). Prurigo nodularis, approve.</p> <p>Recurrent aphthous ulcers or aphthous stomatitis, approve if the patient has tried two other therapies (eg, topical or intralesional corticosteroids, systemic corticosteroids, topical anesthetics/analgesics [eg, benzocaine lozenges], antimicrobial mouthwashes [eg, tetracycline], acyclovir, colchicine). AIDS Related Kaposi's Sarcoma-approve if the patient has tried one regimen and has relapsed or refractory disease.</p>
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	<p>Discoid lupus erythematosus or cutaneous lupus erythematosus, Myelofibrosis, Prurigo nodularis, Recurrent aphthous ulcers or aphthous stomatitis, Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma, Systemic Light Chain Amyloidosis, AIDS related Kaposi's Sarcoma, Castleman's Disease (relapsed/refractory or progressive).</p>

TIBSOVO

Products Affected

- Tibsovo

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, IDH1 Status
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	AML- approve if the disease is isocitrate dehydrogenase-1 (IDH1) mutation positive, as detected by an approved test.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

TOPICAL AGENTS FOR ATOPIC DERMATITIS

Products Affected

- tacrolimus topical

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	Authorize use in patients who have tried a prescription strength topical corticosteroid (brand or generic) for the current condition. Dermatologic condition on or around the eyes, eyelids, axilla, or genitalia, authorize use without a trial of a prescription strength topical corticosteroid.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TOPICAL RETINOID PRODUCTS

Products Affected

- tretinoin topical

PA Criteria	Criteria Details
Exclusion Criteria	Coverage is not provided for cosmetic use.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TOPICAL TESTOSTERONE PRODUCTS

Products Affected

- testosterone transdermal gel in metered-dose pump 20.25 mg/1.25 gram (1.62 %)
- testosterone transdermal gel in packet 1 % (25 mg/2.5gram), 1.62 % (20.25 mg/1.25 gram), 1.62 % (40.5 mg/2.5 gram)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of primary hypogonadism (congenital or acquired) in males. Diagnosis of secondary (hypogonadotropic) hypogonadism (congenital or acquired) in males. Hypogonadism (primary or secondary) in males, serum testosterone level. [Man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression.]
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	Hypogonadism (primary or secondary) in males - initial therapy, approve if all of the following criteria are met: 1) patient has persistent signs and symptoms of androgen deficiency (pre-treatment) [eg, depressed mood, decreased energy, progressive decrease in muscle mass, osteoporosis, loss of libido, AND 2) patient has had two pre-treatment serum testosterone (total or available) measurements, each taken in the morning on two separate days, AND 3) the two serum testosterone levels are both low, as defined by the normal laboratory reference values. hypogonadism has been confirmed by a low for age serum testosterone (total or free) level defined by the normal laboratory reference values. Hypogonadism (primary or secondary) in males - continuing therapy, approve if the patient meets all of the following criteria: 1) patient has persistent signs and symptoms of androgen deficiency (pre-treatment) AND 2) patient had at least one pre-treatment serum testosterone level that was low. [Note: male is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression.]
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off-Label Uses	N/A

TOPIRAMATE/ZONISAMIDE

Products Affected

- topiramate oral capsule, sprinkle
- topiramate oral tablet
- zonisamide

PA Criteria	Criteria Details
Exclusion Criteria	Coverage is not provided for weight loss or smoking cessation.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TRANSDERMAL FENTANYL

Products Affected

- fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 25 mcg/hr, 50 mcg/hr, 75 mcg/hr

PA Criteria	Criteria Details
Exclusion Criteria	Acute (i.e., non-chronic) pain
Required Medical Information	Pain type (chronic vs acute), prior pain medications/therapies tried, concurrent pain medications/therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	For pain severe enough to require daily, around-the-clock, long-term opioid treatment, approve if all of the following criteria are met: 1) patient is not opioid naive, AND 2) non-opioid therapies have been optimized and are being used in conjunction with opioid therapy according to the prescribing physician, AND 3) the prescribing physician has checked the patient's history of controlled substance prescriptions using state prescription drug monitoring program (PDMP), AND 4) the prescribing physician has discussed risks (eg, addiction, overdose) and realistic benefits of opioid therapy with the patient, AND 5) according to the prescriber physician there is a treatment plan (including goals for pain and function) in place and reassessments are scheduled at regular intervals. Patients with cancer, sickle cell disease, in hospice or who reside in a long term care facility are not required to meet above criteria. Clinical criteria incorporated into the quantity limit edits for all oral long-acting opioids (including transdermal fentanyl products) require confirmation that the indication is intractable pain (ie, FDA labeled use) prior to reviewing for quantity exception.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

TRANSMUCOSAL FENTANYL DRUGS

Products Affected

- fentanyl citrate buccal lozenge on a handle

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 12 months.
Other Criteria	For breakthrough pain in patients with cancer if patient is unable to swallow, has dysphagia, esophagitis, mucositis, or uncontrollable nausea/vomiting OR patient is unable to take 2 other short-acting narcotics (eg, oxycodone, morphine sulfate, hydromorphone, etc) secondary to allergy or severe adverse events AND patient is on or will be on a long-acting narcotic (eg, Duragesic), or the patient is on intravenous, subcutaneous, or spinal (intrathecal, epidural) narcotics (eg, morphine sulfate, hydromorphone, fentanyl citrate). Clinical criteria incorporated into the quantity limit edits for all transmucosal fentanyl drugs require confirmation that the indication is breakthrough cancer pain (ie, FDA labeled use) prior to reviewing for quantity exception.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

TUKYSA

Products Affected

- Tukysa oral tablet 150 mg, 50 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years
Other Criteria	Breast Cancer-approve if the patient has advanced unresectable or metastatic human epidermal growth factor receptor 2 (HER2)-positive disease, has received at least one prior anti-HER2-based regimen in the metastatic setting and Tukysa is used in combination with trastuzumab and capecitabine.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

TYKERB

Products Affected

- Tykerb

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Tykerb is being used. Metastatic breast cancer, HER2 status or hormone receptor (HR) status.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	HER2-positive advanced or metastatic breast cancer, approve if Tykerb will be used in combination with Xeloda or Herceptin and the patient has received prior therapy with Herceptin. HER2-positive HR positive metastatic breast cancer, approve if the patient is a man receiving a GnRH agonist, a premenopausal or perimenopausal woman receiving ovarian suppression/ablation with a GnRH agonist, surgical bilateral oophorectomy or ovarian radiation or a postmenopausal woman and Tykerb will be used in combination with an aromatase inhibitor, that is letrozole (Femara), anastrozole, or exemestane. In this criteria, man/woman is defined as an individual with the biological traits of a man/woman, regardless of the individual's gender identity or expression. Colon or rectal cancer-approve if the patient has unresectable advanced or metastatic disease that is human epidermal receptor 2 (HER2) amplified and with wild-type RAS and the medication is used as subsequent therapy in combination with trastuzumab and the patient has not been previously treated with a HER2-inhibitor.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Bone cancer-chordoma, colon or rectal cancer

TYMLOS

Products Affected

- Tymlos

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other medications for osteoporosis (eg, denosumab [Prolia], bisphosphonates, calcitonin nasal spray [Fortical], Forteo), except calcium and Vitamin D. Previous use of Tymlos and/or Forteo for a combined total no greater than 2 years duration during a patient's lifetime.
Required Medical Information	Previous medications tried, renal function
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 2 years of therapy over a patient's lifetime
Other Criteria	Treatment of PMO, approve if the patient meets ONE of the following criteria: patient has tried one oral bisphosphonate or cannot take an oral bisphosphonate because the patient cannot swallow or has difficulty swallowing or patient cannot remain in an upright position post oral bisphosphonate administration or patient has a pre-existing GI medical condition (eg, patient with esophageal lesions, esophageal ulcers, or abnormalities of the esophagus that delay esophageal emptying [stricture, achalasia]), OR patient has tried an IV bisphosphonate (ibandronate or zoledronic acid), OR patient has severe renal impairment or CKD, OR patient has had an osteoporotic fracture or fragility fracture
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

UPTRAVI

Products Affected

- Uptravi

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Confirmation of right heart catheterization, medication history.
Age Restrictions	N/A
Prescriber Restrictions	PAH must be prescribed by, or in consultation with, a cardiologist or a pulmonologist.
Coverage Duration	1 year
Other Criteria	Must have PAH (WHO Group 1) and had a right heart catheterization to confirm the diagnosis of PAH (WHO Group 1). Patient new to Uptravi therapy must meet a) OR b): a) tried one or is currently taking one oral therapy for PAH for 30 days, unless patient has experienced treatment failure, intolerance, or oral therapy is contraindicated: PDE5 inhibitor (eg, sildenafil, Revatio), endothelin receptor antagonist (ERA) [eg, Tracleer, Letairis or Opsumit], or Adempas, OR b) receiving or has received in the past one prostacyclin therapy for PAH (eg, Orenitram, Ventavis, or epoprostenol injection).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

VALCHLOR

Products Affected

- Valchlor

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

VALTOCO

Products Affected

- Valtoco

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other medications used at the same time, prior therapies
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	1 year
Other Criteria	Intermittent Episodes of Frequent Seizure Activity (i.e., seizure clusters, acute repetitive seizures)-approve if the patient is currently receiving maintenance antiepileptic medication(s).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

VENCLEXTA

Products Affected

- Venclexta oral tablet 10 mg, 100 mg, 50 mg
- Venclexta Starting Pack

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	CLL with or without 17p deletion - approve. SLL-approve. Mantle Cell Lymphoma-approve if the patient has tried one prior therapy. AML-approve if the patient is using Venclexta in combination with either azacitidine, decitabine, or cytarabine.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Mantle Cell Lymphoma

VERZENIO

Products Affected

- Verzenio

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	HR status, HER2 status, previous medications/therapies tried, concomitant therapy, menopausal status
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years

PA Criteria	Criteria Details
Other Criteria	<p>Breast cancer - approve advanced or metastatic hormone receptor positive (HR+) [i.e., estrogen receptor positive (ER+) and/or progesterone receptor positive (PR+)] disease, and HER2-negative breast cancer in patients who have not had disease progression while on Kisqali, Ibrance or Verzenio when the pt meets ONE of the following</p> <ol style="list-style-type: none"> 1. Pt is postmenopausal and Verzenio will be used in combination with anastrozole, exemestane, or letrozole 2. pt is premenopausal or perimenopausal and is receiving ovarian suppression/ablation with GnRH agonist, or has had surgical bilateral oophorectomy, or ovarian irradiation AND Verzenio will be used in combination with anastrozole, exemestane, or letrozole 3. Patient is postmenopausal and meets the following conditions: Verzenio will be used in combination with Faslodex. 4. patient is premenopausal or perimenopausal and meets the following conditions: The patient is receiving ovarian suppression/ablation with a gonadotropin-releasing hormone (GnRH) agonist, or has had surgical bilateral oophorectomy or ovarian irradiation AND Verzenio will be used in combination with Faslodex 5. patient is postmenopausal, premenopausal/perimenopausal (patient is receiving ovarian suppression/ablation with GnRH agonist or has had surgical bilateral oophorectomy or ovarian irradiation) or a man (a man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression) and meets the following conditions: Verzenio will be used as monotherapy AND patient's breast cancer has progressed on at least one prior endocrine therapy (e.g., anastrozole, exemestane, letrozole, tamoxifen, Fareston, exemestane plus Afinitor, Faslodex, Afinitor plus Faslodex or tamoxifen, megestrol acetate, fluoxymesterone, ethinyl estradiol) AND patient has tried chemotherapy for metastatic breast cancer. 6. pt is a man who is receiving GnRH agonist AND Verzenio will be used in combination with anastrozole, exemestane or letrozole 7. Patient is a man and Verzenio will be used in combination with Faslodex
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Men with breast cancer

VITRAKVI

Products Affected

- Vitrakvi oral capsule 100 mg, 25 mg
- Vitrakvi oral solution

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, NTRK gene fusion status
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Solid tumors - approve if the tumor has a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation AND the tumor is metastatic or surgical resection of tumor will likely result in severe morbidity AND there are no satisfactory alternative treatments or the patient has disease progression following treatment.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

VIZIMPRO

Products Affected

- Vizimpro

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, EGFR status, exon deletions or substitutions
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Metastatic-NSCLC-Epidermal Growth Factor Receptor (EGFR) mutation positive AND has epidermal growth factor receptor (EGFR) exon 19 deletion as detected by an approved test OR exon 21 (L858R) substitution mutations as detected by an approved test.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

VOTRIENT

Products Affected

- Votrient

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	Soft tissue sarcoma other than GIST [angiosarcoma, Pleomorphic rhabdomyosarcoma, retroperitoneal/intra-abdominal soft tissue sarcoma that is unresectable or progressive, soft tissue sarcoma of the extremity/superficial trunk or head/neck, including synovial sarcoma, or solitary fibrous tumor/hemangiopericytoma or alveolar soft part sarcoma], approve. Differentiated (ie, papillary, follicular, Hurthle) thyroid carcinoma, approve if the patient is refractory to radioactive iodine therapy. Uterine sarcoma, approve if the patient has recurrent, advanced or metastatic disease. Renal Cell Carcinoma, Clear Cell or non-Clear Cell histology-approved if the patient has relapsed or stage IV disease. Ovarian Cancer (ie, Epithelial Ovarian, Fallopian Tube, or Primary Peritoneal Cancer) - approve if the patient has persistent or recurrent disease. GIST - approve if the patient has tried TWO of the following: Gleevec, Sutent, or Strivarga. Medullary Thyroid Carcinoma, approve if the patient has tried vandetanib (Caprelsa) or cabozantinib (Cometriq).
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Differentiated (ie, papillary, follicular, Hurthle cell) thyroid carcinoma. Uterine sarcoma, Epithelial Ovarian, Fallopian Tube, or Primary Peritoneal Cancer, Gastrointestinal Stromal Tumor (GIST), Medullary thyroid carcinoma.

XALKORI

Products Affected

- Xalkori

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For the FDA-approved indication of NSCLC for patients new to therapy, ALK status, high level MET amplification status, MET Exon 14 skipping mutation, and ROS1 rearrangement required. For soft tissue sarcoma IMT, ALK translocation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	NSCLC, patient new to therapy must be ALK-positive, have high level MET amplification, have MET Exon 14 skipping mutation, or have ROS1 rearrangement for approval. For IMT, patient new to therapy must have ALK translocation for approval.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Soft tissue sarcoma Inflammatory Myofibroblastic Tumor (IMT) with ALK translocation, Plus peripheral T-Cell Lymphoma - Anaplastic Large Cell Lymphoma (ALCL), ALK PositivePlus NSCLC with high level MET amplification or MET Exon 14 skipping mutation.

XCOPRI

Products Affected

- Xcopri
- Xcopri Maintenance Pack
- Xcopri Titration Pack

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, medication history
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Partial onset seizures-new starts-approve if the patient has tried one other anticonvulsant therapy (eg, carbamazepine, divalproex sodium, lamotrigine, levetiracetam, oxcarbazepine, phenytoin, topiramate, valproic acid).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

XELJANZ

Products Affected

- Xeljanz
- Xeljanz XR oral tablet extended release 24 hr 11 mg, 22 mg

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with a biologic or with a Targeted Synthetic DMARD for an inflammatory condition (eg, tocilizumab, anakinra, abatacept, rituximab, certolizumab pegol, etanercept, adalimumab, infliximab, golimumab). Concurrent use with potent immunosuppressants that are not methotrexate (MTX) [eg, azathioprine, tacrolimus, cyclosporine, mycophenolate mofetil].
Required Medical Information	Diagnosis, concurrent medications, previous drugs tried.
Age Restrictions	N/A
Prescriber Restrictions	RA, prescr/consult w/rheum. PsA-prescr/consult w/rheum or a derm. UC-prescr/consult w/gastro.
Coverage Duration	PsA/RA-3 months initial, UC-16 weeks initial, All diagnoses-3 years cont.
Other Criteria	RA/PsA initial, patient has tried one conventional synthetic DMARD for at least 3 months (note: patients who have already had a 3-month trial of a biologic for RA are not required to step back and try a conventional synthetic DMARD). UC-Approve if the patient has had a trial of at least ONE tumor necrosis factor inhibitor for ulcerative colitis. Continuation Therapy - Patient must have responded, as determined by the prescriber.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

XENAZINE

Products Affected

- tetrabenazine oral tablet 12.5 mg, 25 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	For treatment of chorea associated with Huntington's disease, Tourette syndrome or related tic disorders, hyperkinetic dystonia, or hemiballism, must be prescribed by or after consultation with a neurologist. For TD, must be prescribed by or after consultation with a neurologist or psychiatrist.
Coverage Duration	Authorization will be for 1 year.
Other Criteria	N/A
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Tardive dyskinesia (TD). Tourette syndrome and related tic disorders. Hyperkinetic dystonia. Hemiballism.

XERMELO

Products Affected

- Xermelo

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapy, concomitant therapy
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Initial therapy - approve if the patient meets ALL of the following criteria: 1) patient has been on long-acting somatostatin analog (SSA) therapy (eg, Somatuline Depot [lanreotide for injection]), AND 2) while on long-acting SSA therapy (prior to starting Xermelo), the patient continues to have at least four bowel movements per day, AND 3) Xermelo will be used concomitantly with a long-acting SSA therapy. Continuation therapy - approve if the patient is continuing to take Xermelo concomitantly with a long-acting SSA therapy for carcinoid syndrome diarrhea.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

XIFAXAN

Products Affected

- Xifaxan oral tablet 200 mg, 550 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Traveler's diarrhea - 12 years of age or older. Hepatic encephalopathy, irritable bowel syndrome - 18 years of age or older.
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XOLAIR

Products Affected

- Xolair subcutaneous recon soln
- Xolair subcutaneous syringe 150 mg/mL, 75 mg/0.5 mL

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with an Interleukin (IL) Antagonist Monoclonal Antibody
Required Medical Information	Moderate to severe persistent asthma and SAR/PAR, baseline IgE level of at least 30 IU/mL. For asthma, patient has a baseline positive skin test or in vitro testing (ie, a blood test for allergen-specific IgE antibodies such as an enzyme-linked immunoabsorbant assay (eg, immunoCAP, ELISA) or the RAST) for 1 or more perennial aeroallergens (eg, house dust mite, animal dander [dog, cat], cockroach, feathers, mold spores) and/or for 1 or more seasonal aeroallergens (grass, pollen, weeds). For SAR/PAR, patient has baseline positive skin testing (eg, grass, tree, or weed pollen, mold spores, house dust mite, animal dander, cockroach) and/or baseline positive in vitro testing (ie, a blood test for allergen-specific IgE antibodies) for one or more relevant allergens (eg, grass, tree, or weed pollen, mold spores, house dust mite, animal dander, cockroach). CIU - must have urticaria for more than 6 weeks (prior to treatment with Xolair), with symptoms present more than 3 days/wk despite daily non-sedating H1-antihistamine therapy (e.g., cetirizine, desloratadine, fexofenadine, levocetirizine, loratadine).
Age Restrictions	Moderate to severe persistent asthma-6 years and older. All other diagnoses-12 years and older
Prescriber Restrictions	Moderate to severe persistent asthma/SAR/PAR if prescribed by, or in consultation with an allergist, immunologist, or pulmonologist. CIU if prescribed by or in consultation with an allergist, immunologist, or dermatologist.
Coverage Duration	Initial tx 4 months, continued tx 12 months

PA Criteria	Criteria Details
Other Criteria	<p>Moderate to severe persistent asthma approve if pt meets criteria 1 and 2: 1) pt has received at least 3 months of combination therapy with an inhaled corticosteroid and at least one the following: long-acting beta-agonist (LABA), long-acting muscarinic antagonist (LAMA), leukotriene receptor antagonist, or theophylline, and 2) patient's asthma is uncontrolled or was uncontrolled prior to receiving any Xolair or anti-IL-4/13 therapy (Dupixent) therapy as defined by ONE of the following (a, b, c, d, or e): a) The patient experienced two or more asthma exacerbations requiring treatment with systemic corticosteroids in the previous year OR b) The patient experienced one or more asthma exacerbation requiring hospitalization or an Emergency Department (ED) visit in the previous year OR c) Patient has a forced expiratory volume in 1 second (FEV1) less than 80% predicted OR d) Patient has an FEV1/forced vital capacity (FVC) less than 0.80 OR e) The patient's asthma worsens upon tapering of oral corticosteroid therapy NOTE: An exception to the requirement for a trial of one additional asthma controller/maintenance medication can be made if the patient has already received anti-IL-4/13 therapy (Dupixent) used concomitantly with an ICS for at least 3 consecutive months. For continued Tx for asthma - patient has responded to therapy as determined by the prescribing physician and continues to receive therapy with one inhaled corticosteroid or inhaled corticosteroid containing combination product. SAR/PAR - approve if pt has tried concurrent therapy with at least one drug from two of the following classes: an oral non-sedating or low-sedating antihistamine, a nasal antihistamine, a nasal corticosteroid, or montelukast. For continued tx SAR/PAR - pt must have responded to therapy as determined by the prescribing physician. For CIU cont tx - must have responded to therapy as determined by the prescribing physician.</p>
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Seasonal or perennial allergic rhinitis (SAR or PAR).

XOSPATA

Products Affected

- Xospata

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, FLT3-mutation status
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	AML - approve if the patient has relapsed or refractory disease AND the disease is FLT3-mutation positive as detected by an approved test.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

XPOVIO

Products Affected

- Xpovio oral tablet 100 mg/week (20 mg x 5), 60 mg/week (20 mg x 3), 80 mg/week (20 mg x 4), 80mg twice week (160 mg/week)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Multiple Myeloma-Approve if the if the patient meets ALL of the following (A, B, and C): A) The patient has tried at least two proteasome inhibitors. Note: Examples include Velcade (bortezomib injection), Kyprolis (carfilzomib infusion), Ninlaro (ixazomib capsules) AND B) The patient has tried at least two immunomodulatory drugs. Note: Examples include Revlimid (lenalidomide capsules), Pomalyst (pomalidomide capsules), Thalomid (thalidomide capsules) AND C) The patient has tried an anti-CD38 monoclonal antibody.Note: For example, Darzalex (daratumumab infusion). Diffuse large B-cell lymphoma-approve if the patient has been treated with at least two prior systemic therapies.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

XTANDI

Products Affected

- Xtandi

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis for which Xtandi is being used. For metastatic castration-resistant prostate cancer, prior therapies tried.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

XYREM

Products Affected

- Xyrem

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Medication history
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by a sleep specialist physician or a Neurologist
Coverage Duration	12 months.
Other Criteria	For Excessive daytime sleepiness (EDS) in patients with narcolepsy - approve if the patient has tried one CNS stimulant (e.g., methylphenidate, dexamethylphenidate, dextroamphetamine), modafinil, or armodafinil and narcolepsy has been confirmed with polysomnography and a multiple sleep latency test (MSLT). Cataplexy treatment in patients with narcolepsy-approve if narcolepsy has been confirmed with polysomnography and a multiple sleep latency test (MSLT).
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ZEJULA

Products Affected

- Zejula

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Ovarian, fallopian tube, or primary peritoneal cancer, maintenance therapy - approve if the patient has recurrent disease and is in complete or partial response after platinum-based chemotherapy regimen. Ovarian, fallopian tube, or primary peritoneal cancer, treatment-approve per label if the patient has tried at least three prior chemotherapy regimens and has homologous recombination deficiency (HRD)-positive disease as confirmed by an approved test.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ZELBORAF

Products Affected

- Zelboraf

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	BRAFV600 mutation status required.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	Melanoma, patient new to therapy must have BRAFV600 mutation for approval AND have unresectable, advanced or metastatic melanoma. HCL - must have relapsed or refractory disease AND tried at least one other therapy for hairy cell leukemia. Thyroid Cancer-patient has disease that is refractory to radioactive iodine therapy. Colon or Rectal cancer-approve if the patient meets the following (A, B, and C): A) The patient has BRAF V600E mutation-positive disease AND B) The patient has previously received a chemotherapy regimen for colon or rectal cancer AND C) The agent is prescribed as part of a combination regimen for colon or rectal cancer. Erdheim-Chester disease, in patients with the BRAF V600 mutation-approve.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Patients with Hairy Cell Leukemia, Non-Small Cell Lung Cancer (NSCLC) with BRAF V600E Mutation, Differentiated thyroid carcinoma (i.e., papillary, follicular, or Hurthle cell) with BRAF-positive disease, patients with colon or rectal cancer

ZEPATIER

Products Affected

- Zepatier

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with other direct acting antivirals, excluding ribavirin or Sovaldi.
Required Medical Information	Genotype, prior medication therapy, concurrent medications, NS5A polymorphism status, prescriber specialty
Age Restrictions	18 years or older
Prescriber Restrictions	Prescribed by or in consultation w/ GI, hepatologist, ID, or liver transplant MD.
Coverage Duration	Will be c/w AASLD guidance and inclusive of treatment already received for the requested drug
Other Criteria	Criteria will be applied consistent with current AASLD/IDSA guidance.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Indications consistent with current AASLD/IDSA guidance

ZOLINZA

Products Affected

- Zolinza

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	CTCL - approve if the patient has progressive, persistent, or recurrent disease on or following treatment with systemic therapies.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ZYDELIG

Products Affected

- Zydelig

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	CLL-approve if the patient has tried one prior therapy. Marginal Zone Lymphoma/Follicular B-Cell Non-Hodgkin Lymphoma/SLL - approve if the patient has tried two prior therapies.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Marginal Zone Lymphoma

ZYKADIA

Products Affected

- Zykadia oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Must have metastatic NSCLC that is anaplastic lymphoma kinase (ALK)-positive as detected by an approved test or ROS1 Rearrangement. IMT - ALK Translocation status.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.
Other Criteria	N/A
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Soft Tissue Sarcoma Inflammatory Myofibroblastic Tumor (IMT) with ALK Translocation. Patients with NSCLC with ROS1 Rearrangement-First-line therapy.

ZYPREXA RELPREVV

Products Affected

- Zyprexa Relprevv intramuscular suspension for reconstitution 210 mg

PA Criteria	Criteria Details
Exclusion Criteria	Dementia-related psychosis
Required Medical Information	N/A
Age Restrictions	Patients aged less than 65 years, approve. Patients aged 65 years and older, other criteria apply.
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

ZYTIGA

Products Affected

- abiraterone

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Authorization will be for 3 years.

PA Criteria	Criteria Details
Other Criteria	<p>Prostate Cancer-Metastatic, Castration-Resistant (mCRPC) and Metastatic, Castration-Sensitive (mCSPC), high risk-Approve if abiraterone is being used in combination with prednisone. Prostate Cancer - Regional Risk Group or Locally Advanced. Approve if the patient meets all of the following criteria (A, B, and C): A) abiraterone is used in combination with prednisone AND B) Patient has regional lymph node metastases and no distant metastases AND C) Patient meets one of the following criteria (i or ii): i. abiraterone with prednisone is used in combination with gonadotropin-releasing hormone (GnRH) analog (e.g., Lupron [leuprolide acetate for injection], Lupron Depot [leuprolide acetate for depot suspension], Trelstar [triptorelin pamoate for injectable suspension], Zoladex [goserelin acetate implant], Vantas [histrelin acetate subcutaneous implant], Firmagon) OR ii. Patient has had an orchiectomy. Prostate Cancer - Metastatic (Castration-Resistant or Castration-Sensitive), Post-External Beam Radiation Therapy (EBRT)-Approve if the patient meets all of the following criteria (A, B, C, D, and E): A) abiraterone is used in combination with prednisone AND B) Patient meets one of the following criteria (i or ii): i. abiraterone with prednisone is used in combination with gonadotropin-releasing hormone (GnRH) analog (e.g., Lupron [leuprolide acetate for injection], Lupron Depot [leuprolide acetate for depot suspension], Trelstar [triptorelin pamoate for injectable suspension], Zoladex [goserelin acetate implant], Vantas [histrelin acetate subcutaneous implant], Firmagon) OR ii. Patient has had an orchiectomy. C) Patient meets one of the following criteria (i or ii): i. There is an increase in prostate specific antigen (PSA) after EBRT OR ii Patient has had a positive digital rectal exam (DRE) after EBRT AND D) Patient is not a candidate for local therapy AND E) Patient has had a positive bone scan.</p>
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off-Label Uses	Prostate Cancer-Regional Risk Group or Locally Advanced. Plus Prostate Cancer - Metastatic (Castration-Resistant or Castration-Sensitive), Post-External Beam Radiation Therapy (EBRT).

PART B VERSUS PART D

Products Affected

- Abelcet
- acetylcysteine
- Actimmune
- acyclovir sodium intravenous solution
- albuterol sulfate inhalation solution for nebulization 0.63 mg/3 mL, 1.25 mg/3 mL, 2.5 mg /3 mL (0.083 %), 2.5 mg/0.5 mL
- AmBisome
- Aminosyn II 10 %
- Aminosyn II 15 %
- Aminosyn-PF 7 % (sulfite-free)
- amphotericin B
- aprepitant
- azathioprine
- budesonide inhalation suspension for nebulization 0.25 mg/2 mL, 0.5 mg/2 mL, 1 mg/2 mL
- caspofungin
- cromolyn inhalation
- cyclophosphamide oral capsule
- cyclosporine modified
- cyclosporine oral capsule
- dronabinol
- Emend oral suspension for reconstitution
- Engerix-B (PF) intramuscular syringe
- Engerix-B Pediatric (PF) intramuscular syringe
- everolimus (immunosuppressive) oral tablet 0.25 mg, 0.5 mg, 0.75 mg
- Firmagon kit w diluent syringe
- Freamine HBC 6.9 %
- Gengraf oral capsule 100 mg, 25 mg
- Gengraf oral solution
- Hepatamine 8%
- Intralipid intravenous emulsion 20 %
- Intralipid intravenous emulsion 30 %
- Intron A injection
- ipratropium bromide inhalation
- ipratropium-albuterol
- methotrexate sodium
- methotrexate sodium (PF) injection solution
- methylprednisolone oral tablet
- mycophenolate mofetil
- mycophenolate sodium
- Nebupent

- Nephramine 5.4 %
- ondansetron
- ondansetron HCl oral solution
- ondansetron HCl oral tablet
- pentamidine inhalation
- Perforomist
- Plenamine
- Prednisone Intensol
- prednisone oral tablet
- Premasol 10 %
- Prograf oral granules in packet
- Pulmozyme
- Recombivax HB (PF) intramuscular suspension 10 mcg/mL, 40 mcg/mL
- Recombivax HB (PF) intramuscular syringe
- Sandimmune oral solution
- sirolimus
- Synribo
- tacrolimus oral
- tobramycin in 0.225 % NaCl
- Travasol 10 %
- Trelstar intramuscular suspension for reconstitution
- TrophAmine 10 %
- Xatmep
- Xgeva
- Zortress oral tablet 0.25 mg, 0.5 mg, 0.75 mg, 1 mg

Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

This page intentionally left blank

Index

Abelcet.....	227	budesonide inhalation suspension for nebulization 0.25 mg/2 mL, 0.5 mg/2 mL, 1 mg/2 mL.....	227
abiraterone.....	225	buprenorphine HCl sublingual.....	23
acetylcysteine.....	227	buprenorphine transdermal patch weekly 10 mcg/hour, 15 mcg/hour, 20 mcg/hour, 5 mcg/hour.....	102
Actimmune.....	227	Butrans transdermal patch weekly 7.5 mcg/hour.....	102
acyclovir sodium intravenous solution...	227	Bydureon BCise.....	66
acyclovir topical ointment.....	1	Bydureon subcutaneous pen injector.....	66
Adempas.....	2	Byetta subcutaneous pen injector 10 mcg/dose(250 mcg/mL) 2.4 mL, 5 mcg/dose (250 mcg/mL) 1.2 mL.....	66
Afinitor.....	3	Cablivi injection kit.....	25
Afinitor Disperz oral tablet for suspension 2 mg, 3 mg, 5 mg.....	3	Cabometyx oral tablet 20 mg, 40 mg, 60 mg.....	26
Aimovig Autoinjector.....	6	Calquence.....	27
albuterol sulfate inhalation solution for nebulization 0.63 mg/3 mL, 1.25 mg/3 mL, 2.5 mg /3 mL (0.083 %), 2.5 mg/0.5 mL.....	227	Caprelsa oral tablet 100 mg, 300 mg.....	28
Alecensa.....	7	Carbaglu.....	29
Alunbrig oral tablet 180 mg, 30 mg, 90 mg.....	8	caspofungin.....	227
Alunbrig oral tablets,dose pack.....	8	Cayston.....	30
Alyq.....	138	Chemet.....	31
AmBisome.....	227	Chenodal.....	32
ambrisentan.....	100	Cholbam oral capsule 250 mg, 50 mg.....	33
Aminosyn II 10 %.....	227	Cinryze.....	24
Aminosyn II 15 %.....	227	clobazam oral suspension.....	34
Aminosyn-PF 7 % (sulfite-free).....	227	clobazam oral tablet.....	34
amitriptyline.....	78	clomipramine.....	78
amphotericin B.....	227	clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg.....	73
Anadrol-50.....	10	Cometriq oral capsule 100 mg/day(80 mg x1-20 mg x1), 140 mg/day(80 mg x1-20 mg x3), 60 mg/day (20 mg x 3/day).....	35
APOKYN.....	12	Copaxone subcutaneous syringe 20 mg/mL, 40 mg/mL.....	36
aprepitant.....	227	Copiktra.....	37
Arcalyst.....	13	Corlanor oral tablet.....	38
Arikayce.....	14	Cotellic.....	39
Ayvakit.....	15	cromolyn inhalation.....	227
azathioprine.....	227	cyclobenzaprine oral tablet 10 mg, 5 mg..	75
Balversa.....	16	cyclophosphamide oral capsule.....	227
Banzel.....	17	cyclosporine modified.....	227
Benlysta subcutaneous.....	18	cyclosporine oral capsule.....	227
benztropine oral.....	74		
Betaseron subcutaneous kit.....	19		
bexarotene.....	177		
Bosulif oral tablet 100 mg, 400 mg, 500 mg.....	20		
Braftovi oral capsule 75 mg.....	21		
Brukina.....	22		

Cystaran.....	40	fentanyl transdermal patch 72 hour 100	
dalfampridine.....	9	mcg/hr, 12 mcg/hr, 25 mcg/hr, 50	
Daliresp.....	41	mcg/hr, 75 mcg/hr.....	192
Daraprim.....	42	Ferriprox oral tablet.....	60
Daurismo oral tablet 100 mg, 25 mg.....	43	Firazyr.....	61
deferasirox oral tablet, dispersible.....	56	Firdapse.....	62
Demser.....	137	Firmagon kit w diluent syringe.....	227
diazepam oral concentrate.....	73	fluconazole in NaCl (iso-osm)	
diazepam oral solution 5 mg/5 mL (1		intravenous piggyback 200 mg/100 mL,	
mg/mL).....	73	400 mg/200 mL.....	11
diazepam oral tablet.....	73	Freamine HBC 6.9 %.....	227
Doptelet (10 tab pack).....	44	Fycompa oral suspension.....	63
Doptelet (15 tab pack).....	44	Fycompa oral tablet 10 mg, 12 mg, 2 mg,	
Doptelet (30 tab pack).....	44	4 mg, 6 mg, 8 mg.....	63
Dotti.....	79	Gattex 30-Vial.....	64
doxepin oral capsule.....	78	Gengraf oral capsule 100 mg, 25 mg.....	227
doxepin oral concentrate.....	78	Gengraf oral solution.....	227
dronabinol.....	227	Gilotrif.....	65
Dupixent Syringe.....	45	glatiramer subcutaneous syringe 20	
Emend oral suspension for		mg/mL, 40 mg/mL.....	36
reconstitution.....	227	Glatopa subcutaneous syringe 20	
Enbrel.....	48	mg/mL, 40 mg/mL.....	36
Enbrel Mini.....	48	Grastek.....	68
Enbrel SureClick.....	48	Harvoni oral pellets in packet 33.75-150	
Engerix-B (PF) intramuscular syringe....	227	mg, 45-200 mg.....	71
Engerix-B Pediatric (PF) intramuscular		Harvoni oral tablet 90-400 mg.....	71
syringe.....	227	Hepatamine 8%.....	227
Epclusa.....	50	Hetlioz.....	72
Epidiolex.....	51	Humira Pen.....	80
Erivedge.....	53	Humira Pen Crohns-UC-HS Start.....	80
Erleada.....	54	Humira Pen Psor-Uveits-Adol HS.....	80
erlotinib oral tablet 100 mg, 150 mg, 25		Humira subcutaneous syringe kit 10	
mg.....	176	mg/0.2 mL, 20 mg/0.4 mL, 40 mg/0.8 mL	80
Esbriet oral capsule.....	55	Humira(CF) Pedi Crohns Starter	
Esbriet oral tablet 267 mg, 801 mg.....	55	subcutaneous syringe kit 80 mg/0.8 mL,	
estradiol oral.....	79	80 mg/0.8 mL-40 mg/0.4 mL.....	80
estradiol transdermal patch weekly.....	79	Humira(CF) Pen Crohns-UC-HS.....	80
everolimus (antineoplastic).....	3	Humira(CF) Pen Psor-Uv-Adol HS.....	80
everolimus (immunosuppressive) oral		Humira(CF) Pen subcutaneous pen	
tablet 0.25 mg, 0.5 mg, 0.75 mg.....	227	injector kit 40 mg/0.4 mL.....	80
Farydak oral capsule 10 mg, 20 mg.....	57	Humira(CF) subcutaneous syringe kit 10	
Fasenra.....	58	mg/0.1 mL, 20 mg/0.2 mL, 40 mg/0.4 mL	80
fentanyl citrate buccal lozenge on a		hydroxyzine HCl oral tablet.....	76
handle.....	193	Ibrance.....	82
		icatibant.....	61

Iclusig oral tablet 15 mg, 45 mg.....	83	megestrol oral suspension 400 mg/10 mL (40 mg/mL), 625 mg/5 mL (125 mg/mL) 109	
Idhifa.....	84	megestrol oral tablet.....	109
imatinib oral tablet 100 mg, 400 mg.....	85	Mekinist oral tablet 0.5 mg, 2 mg.....	110
Imbruvica oral capsule 140 mg, 70 mg.....	86	Mektovi.....	111
Imbruvica oral tablet.....	86	memantine oral capsule,sprinkle,ER 24hr.....	112
imipramine HCl.....	78	memantine oral solution.....	112
Increlex.....	87	memantine oral tablet.....	112
Inlyta oral tablet 1 mg, 5 mg.....	89	memantine oral tablets,dose pack.....	112
Inrebic.....	90	methadone oral solution 10 mg/5 mL, 5 mg/5 mL.....	102
Intralipid intravenous emulsion 20 %.....	227	methadone oral tablet 10 mg, 5 mg.....	102
Intralipid intravenous emulsion 30 %.....	227	methotrexate sodium.....	227
Intron A injection.....	227	methotrexate sodium (PF) injection solution.....	227
ipratropium bromide inhalation.....	227	methylprednisolone oral tablet.....	227
ipratropium-albuterol.....	227	modafinil oral tablet 100 mg, 200 mg.....	126
Iressa.....	91	morphine oral tablet extended release....	102
Jakafi.....	93	Myalept.....	113
Kalydeco oral granules in packet.....	94	mycophenolate mofetil.....	227
Kalydeco oral tablet.....	94	mycophenolate sodium.....	227
Kisqali Femara Co-Pack oral tablet 200 mg/day(200 mg x 1)-2.5 mg, 400 mg/day(200 mg x 2)-2.5 mg, 600 mg/day(200 mg x 3)-2.5 mg.....	95	Namzaric.....	112
Kisqali oral tablet 200 mg/day (200 mg x 1), 400 mg/day (200 mg x 2), 600 mg/day (200 mg x 3).....	95	Natpara.....	114
Korlym.....	97	Nayzilam.....	115
Kuvan.....	98	Nebupent.....	227
Lenvima oral capsule 10 mg/day (10 mg x 1), 12 mg/day (4 mg x 3), 14 mg/day(10 mg x 1-4 mg x 1), 18 mg/day (10 mg x 1-4 mg x2), 20 mg/day (10 mg x 2), 24 mg/day(10 mg x 2-4 mg x 1), 4 mg, 8 mg/day (4 mg x 2).....	99	Nephramine 5.4 %.....	227
lidocaine topical adhesive patch,medicated 5 %.....	101	Nerlynx.....	116
Lonsurf oral tablet 15-6.14 mg, 20-8.19 mg.....	104	Neulasta subcutaneous syringe.....	117
lorazepam oral concentrate.....	73	Neupogen.....	118
lorazepam oral tablet 0.5 mg, 1 mg, 2 mg	73	Nexavar.....	120
Lorbrena oral tablet 100 mg, 25 mg.....	105	Ninlaro.....	121
Lupron Depot.....	67	Norditropin FlexPro subcutaneous pen injector 10 mg/1.5 mL (6.7 mg/mL), 15 mg/1.5 mL (10 mg/mL), 5 mg/1.5 mL (3.3 mg/mL).....	69
Lupron Depot (3 month).....	67	norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg.....	79
Lupron Depot (4 month).....	67	Northera oral capsule 100 mg, 200 mg, 300 mg.....	122
Lupron Depot (6 Month).....	67	Nubeqa.....	123
Lynparza oral tablet.....	106	Nuedexta.....	124
		Nuplazid oral capsule.....	125
		Nuplazid oral tablet 10 mg.....	125
		Ocaliva.....	127

octreotide acetate injection solution.....	128	Promacta oral tablet 12.5 mg, 25 mg, 50	
Odomzo.....	129	mg, 75 mg.....	144
Ofev.....	130	promethazine oral tablet 25 mg.....	76
ondansetron.....	227	Pulmozyme.....	227
ondansetron HCl oral solution.....	227	pyrimethamine.....	42
ondansetron HCl oral tablet.....	227	Qinlock.....	145
Orencia.....	131	quinine sulfate.....	146
Orencia ClickJect.....	131	Recombivax HB (PF) intramuscular	
Orkambi oral granules in packet.....	132	suspension 10 mcg/mL, 40 mcg/mL.....	227
Orkambi oral tablet.....	132	Recombivax HB (PF) intramuscular	
oxandrolone oral tablet 10 mg, 2.5 mg.....	10	syringe.....	227
Oxervate.....	133	Relistor subcutaneous solution.....	147
oxymorphone oral tablet extended		Relistor subcutaneous syringe.....	147
release 12 hr.....	102	Repatha.....	148
Ozempic subcutaneous pen injector 0.25		Repatha Pushtronex.....	148
mg or 0.5 mg(2 mg/1.5 mL), 1 mg/dose		Repatha SureClick.....	148
(2 mg/1.5 mL).....	66	Retevmo.....	150
Palynziq subcutaneous syringe 10 mg/0.5		Revlimid.....	151
mL, 2.5 mg/0.5 mL, 20 mg/mL.....	134	Rinvoq.....	152
Pegasys ProClick subcutaneous pen		Rozlytrek oral capsule 100 mg, 200 mg..	153
injector 180 mcg/0.5 mL.....	135	Rubraca.....	154
Pegasys subcutaneous solution.....	135	Rydapt.....	155
Pegasys subcutaneous syringe.....	135	Samsca oral tablet 15 mg, 30 mg.....	157
Pemazyre.....	136	Sandimmune oral solution.....	227
pentamidine inhalation.....	227	Signifor.....	158
Perforomist.....	227	sildenafil (Pulmonary Arterial	
phenobarbital oral elixir.....	77	Hypertension) oral suspension for	
phenobarbital oral tablet.....	77	reconstitution 10 mg/mL.....	138
Piqray.....	139	sildenafil (Pulmonary Arterial	
Plenamine.....	227	Hypertension) oral tablet 20 mg.....	138
Pomalyst.....	140	sirolimus.....	227
Prednisone Intensol.....	227	Sirturo oral tablet 100 mg.....	159
prednisone oral tablet.....	227	Skyrizi subcutaneous syringe kit.....	160
Premasol 10 %.....	227	Somatuline Depot.....	161
Privigen.....	92	Somavert.....	162
Procrit injection solution 10,000		Sprycel oral tablet 100 mg, 140 mg, 20	
unit/mL, 2,000 unit/mL, 20,000 unit/mL,		mg, 50 mg, 70 mg, 80 mg.....	163
3,000 unit/mL, 4,000 unit/mL, 40,000		Stelara subcutaneous.....	164
unit/mL.....	52	Stivarga.....	165
Prograf oral granules in packet.....	227	Sutent.....	166
Prolastin-C intravenous recon soln.....	141	Sylatron subcutaneous kit 200 mcg, 300	
Prolia.....	142	mcg.....	167
Promacta oral powder in packet 12.5		Symdeko.....	168
mg, 25 mg.....	144	SymlinPen 120.....	169
		SymlinPen 60.....	169

Sympazan.....	34	Tymlos.....	196
Synribo.....	227	Uptravi.....	197
Tabrecta.....	171	Valchlor.....	198
tacrolimus oral.....	227	Valtoco.....	199
tacrolimus topical.....	187	Venclexta oral tablet 10 mg, 100 mg, 50	
tadalafil (pulm. hypertension).....	138	mg.....	200
Tafinlar.....	173	Venclexta Starting Pack.....	200
Tagrisso.....	174	Verzenio.....	201
Talzenna oral capsule 0.25 mg, 1 mg.....	175	Victoza 3-Pak.....	66
Targretin topical.....	178	vigabatrin.....	156
Tasigna oral capsule 150 mg, 200 mg, 50		Vigadrone.....	156
mg.....	179	Vittrakvi oral capsule 100 mg, 25 mg.....	203
tazarotene.....	180	Vittrakvi oral solution.....	203
Tazorac topical cream 0.05 %.....	180	Vizimpro.....	204
Tazverik.....	181	voriconazole intravenous.....	11
Tecfidera.....	182	Votrient.....	205
teriparatide.....	183	Vyndamax.....	172
testosterone cypionate intramuscular oil		Xalkori.....	206
100 mg/mL, 200 mg/mL, 200 mg/mL (1		Xatmep.....	227
ML).....	88	Xcopri.....	207
testosterone enanthate.....	88	Xcopri Maintenance Pack.....	207
testosterone transdermal gel in metered-		Xcopri Titration Pack.....	207
dose pump 20.25 mg/1.25 gram (1.62 %) 189		Xeljanz.....	208
testosterone transdermal gel in packet 1		Xeljanz XR oral tablet extended release	
% (25 mg/2.5gram), 1.62 % (20.25		24 hr 11 mg, 22 mg.....	208
mg/1.25 gram), 1.62 % (40.5 mg/2.5		Xermelo.....	210
gram).....	189	Xgeva.....	227
tetrabenazine oral tablet 12.5 mg, 25 mg 209		Xifaxan oral tablet 200 mg, 550 mg.....	211
Thalomid oral capsule 100 mg, 150 mg,		Xolair subcutaneous recon soln.....	212
200 mg, 50 mg.....	185	Xolair subcutaneous syringe 150 mg/mL,	
Tibsovo.....	186	75 mg/0.5 mL.....	212
tobramycin in 0.225 % NaCl.....	227	Xospata.....	214
topiramate oral capsule, sprinkle.....	191	Xpovio oral tablet 100 mg/week (20 mg x	
topiramate oral tablet.....	191	5), 60 mg/week (20 mg x 3), 80 mg/week	
Tracleer.....	100	(20 mg x 4), 80mg twice week (160	
Travasol 10 %.....	227	mg/week).....	215
Trelstar intramuscular suspension for		Xtandi.....	216
reconstitution.....	227	Xyrem.....	217
tretinoin topical.....	188	Zejula.....	218
trientine.....	170	Zelboraf.....	219
trimipramine.....	78	Zepatier.....	220
TrophAmine 10 %.....	227	Zolinza.....	221
Trulicity.....	66	zonisamide.....	191
Tukysa oral tablet 150 mg, 50 mg.....	194	Zortress oral tablet 0.25 mg, 0.5 mg, 0.75	
Tykerb.....	195	mg, 1 mg.....	227

ZTlido	101
Zydelig.....	222
Zykadia oral tablet.....	223
Zyprexa Relprevv intramuscular suspension for reconstitution 210 mg.....	224